


LATEST TOARCIAN AMMONOIDS FROM THE NORTH AMERICAN CORDILLERA

by GISELLE K. JAKOBS and PAUL L. SMITH

ABSTRACT. Latest Toarcian (Early Jurassic) strata crop out in southern Alaska, southern Yukon, British Columbia, south-western Alberta, and eastern Oregon. Work in the Queen Charlotte Islands, British Columbia has proved a relatively complete Toarcian sequence which has provided the basis for a North American Toarcian ammonite zonation. The Upper Toarcian Yakounensis Zone is characterized by a diverse ammonite fauna, previously assigned to the Middle Toarcian. Species of *Hammatoceras*, *Dumortieria*, *Sphaerocoeloceras*, *Pleydellia*, *Holcophylloceras*, and *Pseudolioceras* allow correlation with the younger part of the Levesquei Zone of north-west Europe. The new genus *Yakounia* and seven new species (*Yakounia yakounensis*, *Y. pacifica*, *Y. freboldi*, *Y. silvae*, *Pleydellia maudensis*, *P. crassiornata* and *Dumortieria? phantasma*) are introduced, all of which are endemic to western North America. A global regression during the Late Toarcian may have restricted migration between the eastern Pacific and western Tethys resulting in the development of endemic taxa.

WESTERN North America is a tectonically complex area made up of numerous accreted terranes (Text-fig. 1). These are commonly fault-bounded regions and each appears to have a separate and distinct geological history. Palaeomagnetic and palaeontological evidence suggests that some terranes may have undergone significant latitudinal displacement since the Jurassic (Irving *et al.* 1980; Taylor *et al.* 1984; Smith and Tipper 1986; Irving and Yole 1987; Irving and Wynne 1991). Understanding this complex history requires correlation within and between terranes, and between the terranes and the craton. To date, Toarcian strata in western North America have been recognized from Alaska to Oregon, both on the craton (southern Canadian Rocky Mountains) and in the following terranes: Peninsular, Stikine (Stikinia), Quesnel (Quesnellia), Wrangell (Wrangellia), Izee (in Oregon), as well as in several small slivers in south-western British Columbia (Text-figs 1–2).


Previous ammonite workers (Frebold 1957, 1964a, 1964b, 1969, 1976; Frebold *et al.* 1967, 1969; Imlay 1968, 1981; Frebold and Tipper 1970) had difficulty interpreting the Toarcian succession of western North America because of the complex geology of the Cordillera and the lack of stratigraphical sections. They relied on the north-west European zonation and compared the North American fauna with common north-west European taxa. However, work on the Early Jurassic of the Americas has shown consistently that eastern Pacific faunas have closer affinities with Tethyan faunas, contain endemic Pacific species, lack certain European elements, and may have different age ranges for common taxa. For example, a re-assessment of some older collections previously assigned to the Middle Toarcian based on the supposed presence of *Haugia* (Frebold 1976; Imlay 1981), indicates that they are in fact of latest Toarcian age, as had been suggested tentatively by Hall (1987). This latest Toarcian fauna includes *Hammatoceras*, *Sphaerocoeloceras*, *Dumortieria*, *Pleydellia*, and a new genus of the Phymatoceratinae; the genus *Haugia* does not occur along the Pacific rim. Such observations clearly point to the need for a regional zonation. A significant step in this direction has been the discovery of a relatively complete Toarcian succession in the Queen Charlotte Islands, British Columbia (Text-fig. 2; Jakobs 1992; Jakobs *et al.* 1994, 1995). The purpose of this paper is to review the distribution of Late Toarcian rocks in North America, and to describe the latest Toarcian ammonites from successions that form the basis of a North American ammonite zonation.



TEXT-FIG. 1. Generalized terrane map of the Canadian Cordillera.

UPPER TOARCIAN ZONES OF NORTH AMERICA

The two Upper Toarcian ammonite zones detailed below were outlined by Jakobs *et al.* (1995), and have been formally described by Jakobs *et al.* (1994).



TEXT-FIG. 2. Map showing Upper Toarcian localities in western North America. The biostratigraphy of sections 1-8 is shown in Text-figures 7-14. Locality data and faunal listings for Collections 1-52 are available from the British Library as Supplementary Publication No. SUP 14044.

Hillebrandti Zone


This zone (Zone 5 of Jakobs *et al.* 1995) contains a prolific, albeit low diversity fauna that includes *Phymatoceras hillebrandti* Jakobs, 1994, *Podagrosites latescens* (Simpson, 1834), and *Grammoceras thouarsense* (d'Orbigny, 1843). The Hillebrandti Zone correlates roughly with the Thouarsense Zone of north-west Europe, based on the occurrence of *Grammoceras thouarsense* and *Podagrosites latescens*, and correlates with the Copiapense Zone of South America based on the similarity between *Phymatoceras copiapense* (Moericke, 1894) and *P. hillebrandti*.

Middle Jurassic	Baj.	Lower	Graham Island	Yak.
		Upper		
Lower Jurassic	Aalenian	Lower	Phantom Creek	Maude Group
		Toarcian	Yakounensis	
	Hillebrandti			
	Crassicosta			
	Ionica			
	Levisoni			
	Pliensbachian	Kanense	Fannin	
		Carlottense		
		Kunae		
		Freboldi		
		Whiteavesi		
	Sinemurian	Upper	Sandilands	
		Lower		
	Hett.	Upper	Kunga Group	
		Middle		
Lower				

TEXT-FIG. 3. Lower Jurassic stratigraphy of the Queen Charlotte Islands. Hett., Hettangian; Baj., Bajocian; Yak., Yakoun Group.

Yakounensis Zone

This zone (Zones 5a and 6 of Jakobs *et al.* 1995) is widespread in western North America, being recognized in southern Alaska, Stikinia, Wrangellia, Quesnellia, south-western British Columbia, Oregon, and on the craton. The interval is thin and stratigraphical relationships between the different species are sometimes difficult to establish. The diverse fauna includes *Pleydellia maudensis* sp. nov., *Pl. crassiornata* sp. nov., *Pl. aalensis* (Zieten, 1832), *Yakounia yakounensis* gen. et sp. nov., *Y. freboldi* sp. nov., *Y. pacifica* sp. nov., *Y. silvae* sp. nov., *Sphaerocoeloceras brochiiforme* Jaworski, 1926, *Hammatoceras speciosum* Janensch, 1902, *Dumortieria insignisimilis* (Brauns, 1865), *D. raricostata* Géczy, 1967, *D. exacta* Buckman, 1905, *D. cf. dumortieri* Thiollière in Dumortier, 1874, *D.?* cf. *pusilla* Jaworski, 1926, *D.?* *phantasma* sp. nov., *Pseudolioceras compactile* (Simpson, 1855)



TEXT-FIG. 4. Extent of the Toarcian/Aalenian hiatus in the Queen Charlotte Islands. Pl. Pliensbachian; Aal., Aalenian.

and *Holcophylloceras calypso* (d'Orbigny, 1841). It contains several taxa endemic to North America and the eastern Pacific.


Both Frebold (1976) and Imlay (1981) assigned this interval to the Middle Toarcian based on the erroneous recognition of *Haugia* (in fact, specimens of *Pleydellia* and *Yakounia*, a new genus endemic to western North America). A rough correlation can be made with the '*Pleydellia fluitans*' and '*Pleydellia lotharingica*' zones of South America, based on the co-occurrence of *Sphaerocoeloceras brochiiiforme*, *Dumortieria* cf. *pusilla*, and species of *Pleydellia*. Both *Pleydellia lotharingica* (Branco, 1879) and *Phlyseogrammoceras? tenuicostatum* (Jaworski, 1926) are similar to *Pleydellia maudensis* which spans the Yakouensis Zone. According to Poulton and Tipper (1991), the base of the Aalenian Stage in North America is defined by the first appearance of *Tmetoceras scissum* (Benecke, 1865). Although *T. scissum* is abundant above the Yakounensis Zone fauna in east-central Oregon, only a single specimen of *Tmetoceras* has been collected from central Graham Island in the Queen Charlotte Islands, the Aalenian there being more commonly characterized by species of *Planammatoceras*, *Bredyia*, and *Erycitoides*.

OCCURRENCES OF LATE TOARCIAN STRATA IN NORTH AMERICA

Queen Charlotte Islands, British Columbia


Mesozoic strata in the Queen Charlotte Islands include the most complete marine Lower Jurassic depositional sequence in North America (Sutherland Brown 1968; Cameron and Tipper 1985; Jakobs 1990; Tipper *et al.* 1991) (Text-fig. 3). Toarcian strata of the Maude Group are represented by the Fannin, Whiteaves and Phantom Creek formations.

The Whiteaves Formation is a grey-green siltstone, weathering brown-grey, which is recessive and commonly poorly exposed in road and stream cuts in central Graham Island, the Skidegate Inlet area and on Louise Island (Text-fig. 5). Neither bedding nor lamination were observed in the siltstones but sandy layers occur at intervals. The siltstones are rich in pyrite and glauconite; ash layers and buff-weathering, calcareous concretions are common. The contact with the overlying Phantom Creek Formation is conformable on much of Graham Island, whereas a hiatus is probably present on Maude island (Skidegate Inlet) and Louise Island (Text-fig. 4).



TEXT-FIG. 5. Toarcian localities in the Queen Charlotte Islands.

The Phantom Creek Formation is a resistant sandstone unit exposed in stream and road cuts. It is best exposed in central Graham Island where it is 25 m thick. Thin (< 2 m) sequences crop out on Maude Island and Louise Island. The formation can be subdivided into two units, a lower




TEXT-FIG. 6. Location of Upper Toarcian sections in central Graham Island, Queen Charlotte Islands.

Coquinoid Sandstone Member and an upper Belemnite Sandstone Member (Cameron and Tipper 1985); these are separated by an erosional hiatus that increases in magnitude toward the south (Text-fig. 4). In central Graham Island, the two members are conformable at Sections 1–3 (Text-figs 5–6), with a cumulative thickness of over 25 m, but at Sections 5 and 6, the Coquinoid Sandstone Member is absent. At Maude Island, Skidegate Inlet, the Coquinoid Sandstone Member is also absent, and a thin (2–3 m) layer of the Belemnite Sandstone Member separates the Whiteaves Formation from the overlying Yakoun Group. The contact between the Phantom Creek Formation and the overlying Yakoun Group is an angular unconformity, best exposed in central Graham Island.

Southern Alaska

In the Talkeetna Mountains (Text-fig. 2), Early Jurassic strata are represented by the Upper Sinemurian to Upper Toarcian Talkeetna Formation, a thick (4600–5800 m) unit of volcanic and volcaniclastic rocks deposited in a marine to non-marine environment (Imlay 1981). Imlay (1981) identified a Middle Toarcian fauna from the Talkeetna Mountains which he assigned to the



Key for Text-figures 7-14.


north-west European Variabilis Zone. Recent work in the Queen Charlotte Islands has shown that the Alaskan fauna is identical to the Late Toarcian Yakounensis Zone assemblage from the Queen Charlotte Islands and contains *Hammatoceras* sp. indet. (= *Phymatoceras*? sp. of Imlay (1981)), *Yakounia yakounensis* (= *Haugia* cf. *variabilis* (d'Orbigny, 1842) of Imlay (1981)), *Pleydellia maudensis* (= *Haugia* cf. *grandis* Buckman, 1898 and *Haugia* cf. *compressa* Buckman, 1898 of Imlay (1981)), *Pleydellia* sp. indet. (= *Brodieia* cf. *tenuicostatum* var. *nodosa* and probably *Haugia* sp. of Imlay (1981)), and *Pseudolioceras* sp. indet. (Collections 1-9).

A small section of Jurassic strata exposed at Puale Bay (Text-fig. 2) includes the Upper Toarcian to Lower Bajocian Kialagvik Formation, a dark grey to black, sandy siltstone containing some hard, buff sandstone (Imlay 1981). Imlay (1981) assigned a single collection to the Middle Toarcian, Variabilis Zone. The fauna is identical to that from the Talkeetna Mountains and Queen Charlotte Islands and contains *Pleydellia maudensis* (= *Haugia* cf. *compressa* of Imlay (1981)), *Pl.* sp. indet. (= *Haugia* cf. *grandis* of Imlay (1981)), and *Pseudolioceras* sp. indet. (Collection 10).

Northern Stikine Terrane

In the northern part of the Stikine Terrane, sediments of the Lower to Middle Jurassic Laberge Group have yielded Late Toarcian ammonites. Two facies were recognized by Souther (1971): the coarse-grained, near-shore Takwahoni Formation in the south-west; and the argillaceous, basinal Inklin Formation in the north-east. Toarcian fossils occur in the former and possibly in the latter (H. W. Tipper, pers. comm. 1992).

Late Toarcian ammonites identified by Frebold (1964a) from the Whitehorse area, southern Yukon (Text-fig. 2; Collections 12-15) include *Catullocceras*? (probably a *Dumortieria*), *Dumortieria*?, and *Harpoceras*? (probably a *Pseudolioceras*). A poorly preserved specimen of *Yakouonia*? sp. indet. has also been identified (Collection 11). The reported *Grammoceras* aff. *G. fallaciosum* (Bayle, 1878) and *Grammoceras*? *boreale* (Whiteaves, 1889) (Frebold 1964a, p. 17, pl. 7, figs 1-4) are actually Middle Jurassic forms (Poulton and Tipper 1991; D. G. Taylor, pers. comm. 1991). Frebold (1964a, p. 4) mentioned the presence of *Catullocceras*? sp. indet. (= *Dumortieria* cf. *dumortieri*) from the Bennett area (Text-fig. 2; Collections 16-17).



TEXT-FIG. 7. Biostratigraphy of Section 1 (Latitude 53° 25' 20" N, Longitude 132° 15' 45" W) along the Yakoun River, Queen Charlotte Islands.

Central Stikine Terrane

Along the west-central margin of the Stikine Terrane, interbedded sediments and volcanics of the Triassic to Middle Jurassic Hazelton Group have yielded Late Toarcian ammonites. In the Iskut area, the Hazelton Group has been divided into four formations: the volcanogenic Unuk River, Betty Creek, and Mount Dilworth formations, overlain by sediments of the Salmon River Formation (Anderson and Thorkelson 1990). The Salmon River Formation in the Iskut area (Text-fig. 2) has yielded several collections of *Pleydellia* cf. *maudensis* from the Yakounensis Zone (Collections 18–19).

In the Spatsizi area, along the northern margin of the Bowser Basin, sedimentary rocks of the Lower to Middle Jurassic Spatsizi Group have been divided into five formations of which the Melisson Formation is probably Late Toarcian in age (Thomson *et al.* 1986). Isolated localities in the Spatsizi area (Text-fig. 2) have yielded *Pleydellia maudensis*, *Yakounia yakounensis*, *Y.* sp. indet., *Dumortieria?* cf. *pusilla*, *Dumortieria* sp. indet., and *Hammatoceras* sp. indet. (Collections 20–32).

A collection of Toarcian ammonites in volcanoclastic sediments from the Toodoggone area (Text-fig. 2) has yielded *Pleydellia* sp. indet. and *Podagrosites?* sp. indet. (Collection 33).

In the McConnell Creek area (Text-fig. 2), along the eastern margin of the Bowser Basin, Lower to Middle Jurassic sediments of the Hazelton Group have yielded isolated collections of Late Toarcian ammonites (Tipper and Richards 1976) from the Smithers Formation, a unit of interbedded, shallow marine volcanoclastic sediments. Poorly preserved specimens of *Dumortieria* sp. indet. and *Yakounia yakounensis* have been collected from the Yakounensis Zone (Collections 34–35).

Toarcian ammonites occur at isolated localities in the Hazelton area (Text-fig. 2). Two collections in the Hazelton Group yielded *Pleydellia* cf. *maudensis* and *Dumortieria* sp. indet. (Collections 36–37) from the Yakounensis Zone.

Quesnel Terrane


Late Toarcian ammonites including *Pleydellia maudensis*, *Yakounia silvae*, *Polyplectus*, and *Dumortieria?* *phantasma* (Collection 38) have been found in the Manson River area on the Quesnel Terrane (Text-fig. 2).

South-western British Columbia


In the Taseko Lakes area (Text-fig. 2), the Lower to Middle Jurassic (Upper Hettangian to Lower Bajocian) Last Creek Formation rests unconformably on the Tyaughton Group (Umhoefer 1989). The upper part of the poorly exposed Last Creek Formation (Upper Sinemurian to Lower Bajocian) is composed of black, calcareous shales, minor sandstones, and thin ash beds. Toarcian shales of the Taseko Lakes area contain *Dumortieria?* sp. and *Hammatoceras* sp. of the Yakounensis Zone (Collections 39–41).

On the west side of Harrison Lake in south-western British Columbia (Text-fig. 2), the Jurassic Harrison Lake Formation rests unconformably on Triassic rocks (Arthur *et al.* 1993). The lowest two of four members (the Celia Cove and Francis Lake members) have yielded rare ammonites of probable Toarcian age with the highest assemblage containing species of *Dumortieria* indicating the Yakounensis Zone (Collections 42–43).

O'Brien (1987) subdivided the Ladner Group in the Boston Bar area into the Lower Jurassic Boston Bar Formation, a sequence of argillite, siltstone, greywacke and conglomerate, and the Middle Jurassic Dewdney Creek Formation, a sequence of volcanic breccia, lava and argillaceous sediments. This sequence is similar to that found to the south in Manning Park (Text-fig. 2). Toarcian ammonites collected from the Ladner Group of Manning Park were assigned to *Phlyseogrammoceras* aff. *P. dispansiforme* (Wunstorff, 1907) by Frebold (Frebold *et al.* 1969) but are comparable with *Yakounia silvae* from the Yakounensis Zone (Collection 44).



TEXT-FIG. 8. Biostratigraphy of Section 2 (Latitude 53° 25' 05" N, Longitude 132° 15' 30" W) along the Yakoun River, Queen Charlotte Islands.



TEXT-FIG. 9. Biostratigraphy of Section 3 (Latitude 53° 25' 00" N, Longitude 132° 16' 05" W) along the Yakoun River, Queen Charlotte Islands.

Southern Canadian Rocky Mountains

The Jurassic Fernie Formation, deposited on the North American craton, crops out from south-western Alberta to the Peace River area of north-eastern British Columbia. It includes the Toarcian Poker Chip Shale, a laterally extensive black shale, 10–38 m thick, that appears to be overlain conformably by sandstones and siltstones of the Lower Bajocian Rock Creek Member; no Aalenian ammonites have been found (Hall 1984, 1987).

In the southern outcrop area (Text-fig. 2; Collection 45), a Late Toarcian fauna of *Pleydellia maudensis* (= ?Grammocerotinae gen. et sp. indet. of Hall (1987)), *Yakounia silvae* (= ?Grammocerotinae gen. et sp. indet. of Hall (1987)), *Y. yakounensis* (= ?Grammocerotinae gen. et sp. indet. of Hall (1987)), *Sphaerocoeloceras* sp. indet. (= ?Hammatoceratidae gen. et sp. indet. of Hall (1987)), and *Dumortieria? phantasma* (= ?Hildoceratinae gen. et sp. indet. of Hall (1987)) has been collected. Frebold (1976) identified several forms as Middle Toarcian but these are actually Late Toarcian in age and include *Pleydellia maudensis* (= *Haugia* sp. indet. and *Haugia* aff. *H. illustris* (Denckmann, 1887) of Frebold (1976)) and *Yakounia freboldi* (= *Haugia* aff. *H. navis* (Dumortier, 1874) of Frebold (1976)) (Collections 46–51).

BIOSTRATIGRAPHY OF THE MEASURED SECTIONS


Section 1: Yakoun River. This section (Text-figs 5–7) was measured and figured originally by Cameron and Tipper (1985, Section 12) who inadvertently inverted the stratigraphy as a result of misinterpreting a poorly exposed and faulted succession. The section was re-measured during the summers of 1987–90 when lower water levels had increased exposure significantly. Siltstones of the Whiteaves Formation lie conformably beneath sandstones of the Phantom Creek Formation. The boundary between the Hillebrandti and Yakounensis zones is not well documented because of poor exposure. The boundary between the Yakounensis Zone and the Aalenian Stage is similarly not well constrained, although it appears to occur above the contact between the Coquinoid Sandstone and Belemnite Sandstone members (Jakobs 1990). The presence of *Bredya* and *Erycitoides howelli* (White, 1889) at 17 m indicates that such taxa as *Tmetoceras* and *Troitsia* could be expected below. Radiolaria identified by E. S. Carter from concretions at 11.4 m indicate a Late Toarcian age; those at 17 m are Aalenian (Carter and Jakobs 1990).

Section 2: Yakoun River. A sequence similar to Section 1 occurs a few metres upstream (Text-figs 6, 8). The Whiteaves Formation is not exposed, but the Phantom Creek Formation is better exposed. The scarcity of ammonites makes the boundary between the Yakounensis Zone and the Aalenian Stage difficult to determine, but it appears to occur below the contact between the Coquinoid Sandstone and Belemnite Sandstone members of the Phantom Creek Formation.

Section 3: Yakoun River. This section (Text-figs 6, 9), originally measured and figured by Cameron and Tipper (1985, Section 11), was re-measured during the summers of 1987–90. The Whiteaves Formation is overlain by a sandstone unit, and a 10 m thick covered interval separates this sandstone unit from the remainder of the Phantom Creek Formation; a fault is possibly present. The top of the section is faulted.


Section 4: Central Graham Island. This section (Text-figs 6, 10), measured and figured originally by Cameron and Tipper (1985, Section 14), was re-measured during the summers of 1987–90. The Whiteaves Formation appears to be overlain conformably by the Phantom Creek Formation. No fossils were collected from the Whiteaves Formation in this section, but another section nearby yielded *Phymatoceras hillebrandti* just below the contact. The boundary between the Yakounensis Zone and the Aalenian Stage is not well constrained because of poor recovery of fossils.

Sections 5–6: Central Graham Island. These two sections (Text-figs 6, 11–12) were measured at Road 59 in Central Graham Island. Section 5 had been measured and figured previously by




TEXT-FIG. 10. Biostratigraphy of Section 4 (Latitude 53° 23' 35" N, Longitude 132° 15' 30" W) in central Graham Island, Queen Charlotte Islands. Wh., Whiteaves; Hill., Hillebrandti.

Cameron and Tipper (1985, Section 13). The base of both sections is covered. The top of Section 5 is truncated by a fault whereas Section 6 is overlain unconformably by the Yakoun Group. The erosional hiatus between the two members of the Phantom Creek Formation has cut out the Coquinoid Sandstone Member in both sections. A single *Tmetoceras scissum* was collected from a faulted section of the Belemnite Sandstone Member.




TEXT-FIG. 11. Biostratigraphy of Section 5 (Latitude 53° 23' 00" N, Longitude 132° 15' 30" W) in central Graham Island, Queen Charlotte Islands. Whit., Whiteaves; Hill., Hillebrandti.

Sections 7-8: Izee Area, Oregon. South-west of the abandoned town of Izee in east-central Oregon (Text-fig. 2), the basal Warm Springs Member of the Snowshoe Formation has yielded Late Toarcian ammonites (Dickinson and Vigrass 1964; Imlay 1968; Smith 1980). Representatives of this poorly preserved fauna were first described by Imlay (1968) but the stratigraphy at that time




TEXT-FIG. 12. Biostratigraphy of Section 6 (Latitude 53° 23' 00" N, Longitude 132° 15' 30" W) in central Graham Island, Queen Charlotte Islands.

was poorly understood. Stratigraphical sections have now been measured at Sheep Creek Divide (Section 7; Text-fig. 13) and Flat Creek (Section 8; Text-fig. 14) where the recessive siltstones and mudstones of the Warm Springs Member rest with a gradational contact on the resistant, volcanoclastic sandstones of the Hyde Formation. Some of the original identifications of the basal Snowshoe Formation ammonites (Imlay 1968) must now be revised in the light of richer collections



TEXT-FIG. 13. Biostratigraphy of Section 7 (Latitude 44° 03' 17" N, Longitude 119° 25' 54" W) at Sheep Creek Divide, Izee area, Oregon.

and a better understanding of the biostratigraphy. Several species of *Dumortieria* occur together with *Hammatoceras speciosum*, *Hammatoceras* spp. (= *Haugia* spp. of Imlay (1968)), *Pleydellia* spp. (= *Grammoceras* spp. of Imlay (1968)), and *Polyplectus* sp. This association is stratigraphically



TEXT-FIG. 14. Biostratigraphy of Section 8 (Latitude 44° 00' 58" N, Longitude 119° 28' 23" W) at Flat Creek, Izee area, Oregon.

beneath the first occurrence of the Aalenian ammonite *Tmetoceras scissum* (see Imlay 1973) and is assigned to the Yakounensis Zone. Although the Yakounensis Zone correlates in part with the Levesquei Zone of north-west Europe, the ammonite assemblage, and particularly the common occurrence of *Polyplectus* sp., is more typical of Mediterranean successions (Donovan 1958; Géczy 1967; Goy and Martinez 1990).

Detailed locality data for Sections 1 to 8 and Collections 1 to 52 have been deposited with the British Library, Boston Spa, Yorkshire, U.K., as Supplementary Publication No. SUP 14044. The information includes geographical and stratigraphical position, locality numbers, the lithostratigraphical unit sampled, and the fauna present.

SYSTEMATIC PALAEOONTOLOGY

Specimens described and illustrated in this paper are housed at the Geological Survey of Canada in Ottawa (GSC) and the University of British Columbia in Vancouver, Canada (UBC).

Abbreviations are as follows (after Smith 1986): CHW, constrictions per half whorl, counted on adoral half whorl; D, shell diameter at which measurements were made; PRHW, primary ribs per half whorl, counted on the adoral half whorl; U = (UD/D) × 100; UD, umbilical diameter; WH, whorl height; WW, whorl width; WWWH = (WW/WH) × 100. Measurements are in millimetres.

Order AMMONOIDEA von Zittel, 1884
Suborder PHYLLOCERATINA Arkell, 1950
Superfamily PHYLLOCERATACEAE von Zittel, 1884
Family PHYLLOCERATIDAE von Zittel, 1884
Subfamily CALLIPHYLOCERTINAE Spath, 1927
Genus HOLCOPHYLOCERAS Spath, 1927
[= *Salfeldiella* Spath, 1927; *Telegdiceras* Kovács, 1939]

Type species. *Phylloceras mediterraneum* Neumayr, 1871.

Diagnosis. Involute shell with high oval whorl section and plain venter; periodic, acutely sigmoidal or angular constrictions present at all stages of ontogeny; outer half of whorl appears to be ribbed; septal sutures with diphyllic saddles except for first lateral saddle which becomes triphyllic in later forms.

Distribution. *Holcophylloceras* is found world-wide from the Lower Jurassic (Toarcian) to the Cretaceous.

Holcophylloceras calypso (d'Orbigny, 1841)

Plate 5, figures 7–8

*1841 *Ammonites calypso* d'Orbigny, pl. 110, figs 1–3.

1976 *Holcophylloceras calypso* (d'Orbigny); Schlegelmilch, p. 26, pl. 1, fig. 6.

Material. Three specimens collected from calcareous concretions within sandstones of the Phantom Creek Formation, Queen Charlotte Islands (Section 2, loc. 17, talus).

Measurements.	D	UD	U	WH	WW	WWWH	CHW
GSC 107260	27.3	2.8	10.3	14.8	10.5	0.71	4
GSC 107260	—	—	—	8.9	6.4	0.72	—
GSC 107304	—	—	—	6.9	5.7	0.83	—
GSC 107305	17.1	2.3	13.5	8.1	6.9	0.85	3

Description. Involute shell with a high oval whorl section and gently sloping umbilical wall. Ornament consists of about eight sinuous constrictions per whorl. Sutures are diphyllic.

Distribution. *Holcophylloceras calpyso* is found world-wide in the Toarcian.

Suborder AMMONITINA Hyatt, 1889
 Superfamily HILDOCERATACEAE Hyatt, 1867
 Family HILDOCERATIDAE Hyatt, 1867
 Subfamily GRAMMOCERATINAE Buckman, 1904

Genus PLEYDELLIA Buckman, 1899
 [= *Cotteswoldia* Buckman, 1902 (= *Gotteswoldia* Théobald, 1950); *Canavaria* Buckman, 1902
 (non Gemmellaro, 1886) (= *Canavarina* Buckman, 1904); *Walkeria* Buckman, 1902
 (= *Walkericeras* Buckman, 1913)]

Type species. *Pleydellia comata* Buckman, 1899.

Diagnosis. Involute to evolute shell with tall, compressed whorls; whorl section lanceolate to triangular; umbilical shoulder abrupt to moderately rounded; venter carinate with weak ventro-lateral shoulders; ribbing sinuous, prorsiradiate on upper flank, and terminating at ventro-lateral shoulder. Ribs may bifurcate at or near umbilical shoulder, and may fade on inner or outer half of flanks.

Distribution. *Pleydellia* is common in the uppermost Toarcian of Europe, South America and North America.

Pleydellia maudensis sp. nov.

Plate 1, figures 1–2; Plate 2, figures 1–2, 11–12; Text-figure 15A

- 1981 *Haugia* cf. *compressa* Buckman; Imlay, p. 43, pl. 12, figs 3, 7, 9.
 1981 *Haugia* cf. *grandis* Buckman; Imlay, p. 42, pl. 12, figs 4, 10, 15.
 1981 *Brodieia* cf. *B. tenuicostata* var. *nodosa* Jaworski; Imlay, p. 42, pl. 12, fig. 8.
 v 1987 ?*Grammoceras* gen. et sp. indet., Hall, p. 1702, pl. 5, figs A–J, P, W.
 1991 *Phlyseogrammoceras* (?) sp., Tipper *et al.*, pl. 7, fig. 3.
 1992 *Haugia* cf. *compressa* Buckman; Hillebrandt and Smith, pl. 4, fig. 7.
 1992 ?*Haugia* sp. indet., Hillebrandt and Smith, pl. 7, fig. 5.


Derivation of name. After Maude Island in Skidegate Inlet, site of some of the first discoveries of Jurassic fossils in the Queen Charlotte Islands.

Material. About 150 specimens in sandstones and calcareous concretions of the Phantom Creek Formation, Queen Charlotte Islands (Section 1, locs 3, 5, 17; Section 2, locs 3, 7, 9, 12, 17; Section 3, locs 89–92, 94–95, 97–98; Section 4, locs 1–3, 5–6, 8–10, 14; Section 5, locs 2–3, 5; Section 6, locs 1–2). Other specimens from the Upper Toarcian of the Talkeetna Mountains (Text-fig. 2, Collection 2), Puale Bay (Collection 10), the Iskut area (Collections 18–19), the Spatsizi area (Collections 21–22), the Hazelton area (Collection 36), the Manson River area (Collection 38), and the southern Canadian Rocky Mountains (Collections 45–46, 48).

Holotype. GSC 99523 (Pl. 2, figs 11–12) from the lower part of the Phantom Creek Formation (Yakounensis Zone), Yakoun River, Queen Charlotte Islands.

Paratypes. GSC 99519 (Pl. 2, figs 1–2), GSC 99524 (Pl. 1, figs 1–2), GSC 99509, GSC 107267–107277.

TEXT-FIG. 15. Septal suture lines of *Pleydellia maudensis*, *Pleydellia crassiornata* and *Yakounia freboldi* from the Queen Charlotte Islands. A, GSC 99519; Section 4, loc. 14; B, GSC 107347; Section 3, loc. 95; C GSC 107286, Section 4, loc. 4. WH, whorl height.



Measurements.	D	UD	U	WH	WW	WWHH	PRHW
GSC 99523	112.2	40.0	35.7	42.1	23.0	0.55	14
GSC 99523	93.7	31.0	33.1	35.0	24.9	0.71	—
GSC 99523	81.8	24.7	30.2	32.0	19.1	0.60	c. 16
GSC 99519	98.5	28.9	29.3	40.0	21.4	0.54	c. 15
GSC 99519	79.4	22.0	27.7	34.1	19.7	0.58	14
GSC 99519	75.0	20.1	26.8	31.8	—	—	13
GSC 99524	153.0	57.9	37.8	—	—	—	—
GSC 99524	124.0	38.0	30.6	—	—	—	—
GSC 107267	—	—	—	8.3	6.2	0.75	—
GSC 107268	58.5	22.1	37.8	21.6	14.8	0.69	16
GSC 107268	46.1	17.0	36.9	17.0	10.9	0.64	11
GSC 107269	42.9	11.1	25.9	18.4	—	—	—
GSC 107269	37.5	10.0	26.7	—	—	—	—
GSC 107269	34.0	9.0	26.5	15.0	—	—	12
GSC 107270	98.9	32.7	33.1	—	—	—	17
GSC 107271	71.8	22.0	30.6	29.8	18.1	0.61	11
GSC 107271	56.5	17.1	30.3	24.0	14.5	0.60	12
GSC 107272	47.5	12.5	26.3	20.7	10.7	0.52	13
GSC 107273	79.3	25.4	32.0	32.1	—	—	15
GSC 107273	63.7	21.1	33.1	25.1	—	—	15
GSC 107273	58.5	20.0	34.2	22.6	—	—	c. 13
GSC 107274	87.4	29.0	33.2	36.0	—	—	15
GSC 107275	64.7	20.3	31.4	25.2	19.0	0.75	11
GSC 107275	55.2	16.4	29.7	—	—	—	10
GSC 107276	91.1	28.1	30.8	36.2	21.0	0.58	13
GSC 107276	67.3	19.0	28.2	28.4	15.8	0.56	12
GSC 107277	82.6	21.7	26.3	36.1	—	—	13


Diagnosis. Moderately evolute shell; ogival whorl section and flat flanks; umbilical wall gently sloping, becoming undercut on outer whorls; umbilical shoulder gently rounded becoming sharp on outer whorls; venter carinate-sulcate; lateral sulci fade during ontogeny; ribbing sinuous. On inner whorls, primary ribs bifurcate at about one-third flank height. On outer whorls, ribs arise singly and in pairs from small tubercles at umbilical shoulder. Ribs weaken on upper flank. Some ribs bifurcate on upper flank or intercalatory ribs may appear. Ribs terminate at ventro-lateral shoulder.

Description. The holotype, GSC 99523, is a moderately well preserved specimen, septate up to 77.4 mm shell diameter with approximately 190° of body chamber, ending in an incomplete aperture at 111.4 mm shell diameter. One side is slightly distorted. The shell is moderately evolute with an ogival whorl section, flat flanks, and a carinate-sulcate venter. The umbilical wall is gently sloping on the inner whorls becoming steeper with age until, at approximately 82 mm shell diameter, it becomes slightly undercut. The umbilical shoulder is gently rounded on the inner whorls, becoming angular as the umbilical wall steepens. The venter possesses shallow lateral sulci which fade on the outer whorls. On the inner whorls, the primary ribs are slightly prorsiradiate. On the outer whorls, the primary ribs arise from small tubercles at the umbilical shoulder and have a more pronounced prorsiradiate trend. The primary ribs are stronger than the secondary ribs which appear at approximately mid-flank. The paratype, GSC 99519, is a moderately well preserved specimen, septate to 77.1 mm shell diameter with approximately 130° of body chamber, ending in an incomplete aperture at 100.4 mm shell diameter. The phragmocone is damaged on one side and is partially obscured by encrusting bivalves. The body chamber is partially crushed. This specimen illustrates the ribbing on the outer whorls of the phragmocone. From the umbilical tubercles, a primary rib proceeds across the flank and bifurcates at approximately one-quarter to one-third the flank height. At approximately two-thirds to three-quarters the flank height, the ribs may bifurcate again or weak intercalatory ribs may arise. The ribbing fades on the body chamber. The paratype, GSC 99524, is a larger specimen, albeit less well preserved. It is septate to 121.7 mm shell diameter with approximately 190° of body chamber ending in an incomplete aperture at 165 mm shell diameter. One side is poorly preserved and partially obscured by encrusting bivalves. The body chamber is fragmented and parts of the venter have been eroded or broken away. The phragmocone lacks fine detail because of the medium-grained sandstone that forms the matrix. This specimen, whilst having faint ribbing on the outer whorls, maintains the tubercles at the umbilical shoulder and these do not fade but remain prominent. Of the other paratypes, one specimen (GSC 107268) differs slightly in that it is slightly more evolute and has more pronounced umbilical tubercles from which three ribs commonly arise. Two other specimens (GSC 99509 and GSC 107267) are fragments of inner whorls. Both show strong primary ribs which bifurcate at mid-flank. The venter is more strongly carinate-sulcate than on the larger specimens.

Remarks. This form is similar to several Late Toarcian genera including *Pseudolillia*, *Gruneria*, *Phlyseogrammoceras*, as well as *Pleydellia*. The whorl shape and rib pattern on the outer whorls are similar to species of *Pseudolillia* figured by Elmi and Rulleau (1987) but the inner whorls are different. Ribbing on the inner whorls of *Pseudolillia* is much denser and finer, and the ribs tend to bundle or remain single rather than bifurcating as markedly as in *Pleydellia maudensis*. In addition, *Pseudolillia* lacks the small umbilical swellings characteristic of the North American species. *Gruneria* has a similar ribbing pattern in which the primary ribs on the inner whorls bifurcate on the lower to mid-flank; however, the whorl shape is more rounded and ellipsoidal, the ribs are finer and denser, and the outer whorls lack the small umbilical swellings that characterize *Pl. maudensis*. *Phlyseogrammoceras* has a similar ribbing pattern, and a steep to undercut umbilical wall on the outer whorls, but the whorl section is more compressed and the venter is sharper, lacking the ventral sulci of *Pl. maudensis*. Hillebrandt (1987) figured several specimens which he assigned to *Phlyseogrammoceras* (?) *tenuicostatum*, a form similar to the North American species but with a more involute shell. Hall (1987) described specimens from the Fernie Formation (?Grammocerotinae gen. et sp. indet.) which he compared with species of *Pleydellia* such as *Pl. fluitans* (Dumortier, 1874)

EXPLANATION OF PLATE 1

Figs 1–2. *Pleydellia maudensis* sp. nov.; GSC 99524, paratype; Yakounensis Zone, lower part of the Phantom Creek Formation; Queen Charlotte Islands, GSC Loc. No. C-87111, Section 3, loc. 89; × 1.



JAKOBS and SMITH, *Pleydellia maudensis*

and *Pl. lotharingica*. *Pl. fluitans* has coarse, distant, rectiradiate ribs and a bisulcate venter, whereas *Pl. lotharingica* has a narrow venter and an ogival whorl section (Knitter and Ohmert 1983, pl. 3, figs 12–14). Hillebrandt (1987, pl. 13, fig. 7, pl. 14, fig. 8) figured several specimens of *Pl. cf. lotharingica* which have a bisulcate venter on the inner whorls and a similar ribbing pattern, but lack the undercut umbilical shoulder and umbilical swellings on the outer whorls that characterize *Pl. maudensis*. The North American form has a strong carinate-sulcate inner whorl and subdued umbilical swellings that are significantly different from other *Pleydellia*, and a new species designation is warranted.

Pleydellia crassiornata sp. nov.

Plate 3, figures 1–8; Text-figure 15B

Derivation of name. The name refers to the thick ornamentation (Latin *crassus*, thick; *ornatus*, ornament).

Material. About 24 well preserved specimens in sandstones and calcareous concretions of the Phantom Creek Formation (Section 1, locs 5, 17; Section 2, locs 4, 17; Section 3, locs 90, 95; Section 4, locs 6, 14), Queen Charlotte Islands.

Holotype. GSC 99513 (Pl. 3, figs 7–8) from the lower part of the Phantom Creek Formation (middle Yakounensis Zone), Yakoun River, Queen Charlotte Islands.

Paratypes. GSC 99510 (Pl. 3, figs 1–2), GSC 99511 (Pl. 3, figs 3–4), GSC 99512 (Pl. 3, figs 5–6), GSC 107278–107279

Measurements.	D	UD	U	WH	WW	WWWH	PRHW
GSC 99513	48.7	19.4	39.8	—	—	—	—
GSC 99513	38.7	14.5	37.5	—	—	—	9
GSC 99513	—	10.8	—	9.6	7.4	0.77	9
GSC 99511	46.3	15.6	33.7	17.1	9.0	0.53	—
GSC 99511	40.0	12.4	31.0	16.1	9.5	0.59	—
GSC 99511	35.9	11.1	30.9	15.0	8.9	0.59	—
GSC 107278	—	20.3	—	18.1	12.7	0.70	13
GSC 107278	34.5	10.8	31.3	14.0	10.0	0.71	10
GSC 107278	28.3	10.0	35.3	11.5	8.8	0.77	—
GSC 107279	58.3	26.2	44.9	16.5	13.6	0.82	12
GSC 107279	53.1	21.3	40.1	17.2	11.6	0.67	11
GSC 107279	48.3	18.5	38.3	14.1	10.9	0.77	11

EXPLANATION OF PLATE 2


Figs 1–2, 11–12. *Pleydellia maudensis* sp. nov.; Yakounensis Zone, lower part of the Phantom Creek Formation; Queen Charlotte Islands. 1–2, GSC 99519, paratype; GSC Loc. No. C-176555, Section 4, loc. 14. 11–12, GSC 99523, holotype; GSC Loc. no. C-87118, Section 3, loc. 95.

Figs 3–7. *Dumortieria? phantasma* sp. nov.; Yakounensis Zone, lower part of the Phantom Creek Formation; Queen Charlotte Islands. 3–4, GSC 99520, holotype; GSC Loc. No. C-87220, Section 2, loc. 4. 5–6, GSC 99521, paratype; GSC Loc. No. C-93576, Section 2, loc. 4. 7, GSC 99522, paratype; GSC Loc. No. C-87220, Section 2, loc. 4.

Figs 8, 15–16. *Dumortieria? cf. pusilla* Jaworski; Yakounensis Zone, Warm Springs Member of the Snowshoe Formation; Izee area, eastern Oregon. 8, UBC 014; UBC Loc. No. F4-4-E, Section 7, loc. 5. 15, UBC 015; UBC Loc. No. F4-4-E, Section 7, loc. 5. 16, UBC 016; UBC Loc. No. F5-1-4, Collection 52.

Figs 9–10, 13–14. *Dumortieria pusilla* Jaworski; Arroyo Negro Argentina, Locality 22, Section 9, Horizon 6 of Jaworski (1926). 9–10, plaster cast of the paratype. 13–14, plaster cast of the holotype.

Arrows mark start of body chamber. All are $\times 1$.




JAKOBS and SMITH, *Pleydellia* and *Dumortieria*

Diagnosis. Moderately evolute shell; ogival whorl section; umbilical wall gently sloping, becoming steeper during ontogeny; umbilical shoulder gently rounded, becoming more angular during ontogeny; flanks slightly convex; venter carinate with weak lateral sulci on inner whorls; ribbing sinuous; thick, prominent primary ribs split into two or three weaker, secondary ribs at approximately one-third to one-half flank height; secondary ribs terminate at ventro-lateral shoulder; some variation in length of primary ribs.

Description. The holotype, GSC 99513, is a moderately well preserved specimen septate to 13.1 mm umbilical diameter with approximately 180° of body chamber ending in an incomplete aperture at 49.0 mm shell diameter. The body chamber is broken and slightly crushed. The shell is moderately evolute with an ogival whorl section. The umbilical wall is gently sloping and the umbilical shoulder is rounded. The flanks are gently convex and merge into the venter with only weak ventro-lateral shoulders. The venter is carinate, bounded by narrow, smooth strips. The ornament consists of sinuous ribs. The coarse primary ribs arise high on the umbilical wall. On the inner whorls, they are short and appear bullate. The primary ribs split into two or three weaker secondary ribs at approximately one-third to one-half the flank height. The secondary ribs terminate at the ventro-lateral shoulder. The paratype, GSC 99511, is a well preserved specimen septate to 43.1 mm shell diameter with approximately 70° of body chamber ending in an incomplete aperture at 47.5 mm shell diameter. The body chamber is slightly crushed. The shell is moderately evolute with an ogival whorl section and possesses similar shell characteristics to the holotype. The ornament differs slightly by being slightly more subdued, and the primary ribs are longer on the inner whorls and not as bullate. The paratype, GSC 99512, is a moderately well preserved shell, septate to 16.6 mm umbilical diameter with approximately 110° of body chamber ending in an incomplete aperture. Most of the venter on the outer whorl of the phragmocone has been eroded away. This specimen is similar to the previous two but has slightly longer primary ribs on the inner whorls than the holotype. The paratype, GSC 99510, is a moderately well preserved shell with approximately 180° of body chamber. The shell is fragmented and distorted with portions of the phragmocone broken away. This specimen is one of the largest, with a maximum shell diameter of approximately 66 mm. It possesses a similar ribbing pattern to the two previous paratypes. The ornament does not fade on the body chamber. The paratype, GSC 107279, is a well preserved specimen septate to 25.0 mm umbilical diameter and with approximately 190° of body chamber ending in an incomplete aperture. The primary ribs on the inner whorls, while not as bullate as the holotype, are shorter than in the other paratypes. The paratype, GSC 107278, is a moderately well preserved specimen with approximately 180° of body chamber ending in an incomplete aperture. Part of the venter on the body chamber is broken away. It is a smaller specimen than the others, and the whorl section is slightly more depressed. The ornament is similar to the other paratypes.

EXPLANATION OF PLATE 3

- Figs 1–8. *Pleydellia crassiornata* sp. nov.; Yakounensis Zone, lower part of the Phantom Creek Formation; Queen Charlotte Islands. 1–2, GSC 99510, paratype; GSC Loc. No. C-87118, Section 3, loc. 95. 3–4, GSC 99511, paratype; GSC Loc. No. C-87221, Section 2, loc. 17. 5–6, GSC 99512, paratype; GSC Loc. No. C-176555, Section 4, loc. 14. 7–8, GSC 99513, holotype; GSC Loc. No. C-87118, Section 3, loc. 95.
- Figs 9–14. *Pleydellia aalensis* (Zieten); Yakounensis Zone, lower part of the Phantom Creek Formation; Queen Charlotte Islands. 9–10, GSC 99514; GSC Loc. No. C-87107, Section 3, loc. 95. 11–12, GSC 99515; GSC Loc. No. C-87233, Section 1, loc. 17. 13–14, GSC 99516; GSC Loc. No. C-87233, Section 1, loc. 17.
- Fig. 15. *Dumortieria* cf. *exacta* Buckman; UBC 010; Yakounensis Zone, Warm Springs Member of the Snowshoe Formation; Izee area, eastern Oregon, UBC Loc. No. F4-4-E, Section 7, loc. 5.
- Fig. 16. *Dumortieria* cf. *dumortieri* (Thiollière); UBC 011; Yakounensis Zone, Warm Springs Member of the Snowshoe Formation; Izee area, eastern Oregon, UBC Loc. No. F4-4-F, Section 7, loc. 6.
- Fig. 17. *Dumortieria insignisimilis* (Brauns); UBC 012; Yakounensis Zone, Warm Springs Member of the Snowshoe Formation; Izee area, eastern Oregon, UBC Loc. No. F4-4-E, Section 7, loc. 5.
- Fig. 18. *Dumortieria*? cf. *pusilla* Jaworski; GSC 99517; Yakounensis Zone, lower part of the Phantom Creek Formation; Queen Charlotte Islands, GSC Loc. No. C-157740, Section 3, loc. 97.
- Fig. 19. *Dumortieria insignisimilis* (Brauns); GSC 99518; Yakounensis Zone, lower part of the Phantom Creek Formation; Queen Charlotte Islands, GSC Loc. No. C-81736, Section 5, loc. 5.
- Fig. 20. *Dumortieria rariocostata* Géczy; UBC 013; Yakounensis Zone, Warm Springs Member of the Snowshoe Formation; Izee area, eastern Oregon, UBC Loc. No. F4-3-D, Section 7, loc. 4.
- Arrows mark start of body chamber. All are $\times 1$.



JAKOBS and SMITH, *Pleydellia* and *Dumortieria*

Remarks. The shell shape and whorl section are similar to those of *Pleydellia aalensis* as figured by Schlegelmilch (1976, pl. 51, figs 8–9). *Pl. aalensis* encompasses a broad variety of forms and shows wide morphological variability. In general, the style of joined ribs distinguishes *Pl. aalensis* from other species of *Pleydellia* (Buckman 1890, p. 193), some of which have joined ribs but these tend to be fine and bundled, e.g. *Pl. subcompta* (Branco, 1879). The specimens of *Pl. aalensis* illustrated by Buckman (1890, pl. 32, figs 4–10) differ from *Pl. crassiornata* by lacking swollen primaries, by being slightly more involute, and by having a steeper umbilical wall. Géczy (1967) figured several forms of *Pl. aalensis*, creating several new subspecies that are similar to *Pl. crassiornata* but which lack the prominent and pervasive bifurcation and swollen primaries that characterize the North American species. *Pleydellia* from North America that have bifurcate and widely spaced ribbing show some differences in the coarseness of the primary ribs, but this may be intra-specific variation and is probably not sufficient to justify the recognition of two separate species.

Pleydellia crassiornata also shows similarity to the Leioceratinae which, according to Donovan *et al.* (1981, p. 115), evolved from *Pleydellia* during the early Aalenian. Schlegelmilch (1985) figured some species of *Leioceras* and *Staufenia* (pls 9–11) which are similar to *Pl. crassiornata*. *Leioceras* and *Staufenia* have acute venters and simple septal sutures.

Pleydellia aalensis (Zieten, 1832)

Plate 3, figures 9–14

- *1832 *Ammonites aalensis* Zieten, pl. 28, fig. 3.
- 1890 *Grammoceras aalense* (Zieten); Buckman, p. 192, pl. 32, figs 7–8.
- 1976 *Pleydellia aalensis* (Zieten); Schlegelmilch, p. 94, pl. 51, figs 8–9.
- 1983 *Pleydellia aalensis* (Zieten); Knitter and Ohmert, pl. 3, figs 2–3.
- 1990 *Pleydellia aalensis* (Zieten); Goy and Martínez, pl. 4, fig. 10.


Material. 20 specimens in the sandstones and calcareous concretions of the Phantom Creek Formation, Queen Charlotte Islands (Section 1, loc. 17; Section 2, loc. 17; Section 3, locs 89, 95).

<i>Measurements.</i>	D	UD	U	WH	WW	WWHH	PRHW
GSC 99514	30.5	10.6	34.8	11.5	7.6	0.66	10
GSC 99515	24.7	7.5	30.4	10.4	6.7	0.64	8
GSC 99516	38.8	13.5	34.8	14.2	—	—	—
GSC 107280	19.0	6.7	35.3	7.3	5.0	0.68	7

Description. The moderately evolute shell has a high oval whorl section and shallow umbilicus. The umbilical wall and shoulder are gently sloping. Flanks are gently convex and merge into the venter with weak ventrolateral shoulders. The venter is carinate. Ribbing density and strength vary. Primary ribs are slightly prorsiradiate to approximately one-third the flank height where, generally, they bifurcate. Secondary ribs are retriradiate to the ventro-lateral shoulders, then become prorsiradiate and approach, but do not reach, the venter. Intercalary ribs are common.

Remarks. Théobald and Moine (1959) studied *Pl. aalensis* in an attempt to define this variable species more clearly. They concluded that *Pl. aalensis* could be defined as possessing simple ribs that commonly bifurcate near the umbilical shoulder. The ribs are falciform, rounded and fade toward the venter. The whorl section is oval and the flanks are slightly flattened. The U value averages 31.5 but can range from 27 to 36, and WWHH averages 0.58, but can range from 0.50 to 0.66. Two varieties can be distinguished by ribbing density, a coarsely ribbed *aalensis* type and a densely ribbed *tenuicostata* type.

The North American specimens fall within the range of variability of *Pl. aalensis*, although the ribs tend to bifurcate higher on the flanks than in the holotype. However, the specimen figured by Schlegelmilch (1976, pl. 51, fig. 9) has ribs that bifurcate near mid-flank. *Pl. fluens* (Buckman, 1890) has much finer and dense ribbing. *Pl. subcompta* has fine, dense ribbing which bundles in threes or fives at the umbilical shoulder. *Pl. crassiornata* has a coarser ornament and swollen primary ribs.



TEXT-FIG. 16. Septal suture lines of *Dumortiera*, *Dumortiera?*, *Pleydellia* and *Atacamiceras*. A, GSC 107345; Section 2, loc. 4, Queen Charlotte Islands; B, GSC 107346; Section 2, loc. 4, Queen Charlotte Islands; C, Hillebrandt (1987, pl. 8, fig. 2; text-fig. 2a); D, Hillebrandt (1987, pl. 8, fig. 16; text-fig. 2b); E, Hillebrandt (1987, pl. 8, fig. 17; text-fig. 2c); F, Jaworski (1926, pl. 4, fig. 17); G, Jaworski (1926, pl. 4, fig. 22); H, Schlegelmilch (1976, p. 94); I, Schlegelmilch (1976, p. 94); J, Jaworski (1926, pl. 4, fig. 19); K, Schindewolf (1964, p. 293, text-fig. 178); L, Schindewolf (1964, p. 295, text-fig. 179). WH, whorl height.

Distribution. *Pleydellia aalensis* is common in the Upper Toarcian Aalensis Zone of Europe (Fischer 1966; Géczy 1967; Goy and Martínez 1990).

Genus DUMORTIERIA Haug, 1885

[= *Catullocheras* Buckman, 1925; *Dactylogammmites* Buckman, 1925; *Phenakoceras* Maubeuge, 1949 (non Frech, 1902); *Phenakocerites* Maubeuge, 1950]

Type species. *Ammonites levesquei* d'Orbigny, 1844 (subsequent designation by Buckman 1890).

Diagnosis. Moderately evolute planulates with ogival whorl sections; umbilical shoulder abrupt to moderately rounded; venter carinate; ribbing almost rectiradiate, becoming slightly prorsiradiate on upper flanks, terminating at venter; ribbing sparse or dense.

Distribution. *Dumortieria* is found almost world-wide in the Late Toarcian correlatives of the Levesquei Zone (Donovan *et al.* 1981).

Dumortieria insignisimilis (Brauns, 1865)

Plate 3, figures 17, 19

- 1865 *Ammonites insigni similis* Brauns, p. 106, pl. 5, figs 5–7.
 1923 *Dumortieria insignisimilis* (Brauns); Ernst, p. 56, pl. 9, figs 1–3.
 1967 *Dumortieria insignisimilis* (Brauns); Géczy, p. 144, pl. 31, fig. 6.
 1993 *Dumortieria cf. insignisimilis* (Brauns); Arthur *et al.*, p. 32, pl. 1, figs 11–15.

Material. Five specimens from eastern Oregon (Section 7, loc. 5) where the species is best represented. Poorly preserved specimens from the Queen Charlotte Islands (Section 5, loc. 5) and from Harrison Lake (Arthur *et al.* 1993; Text-fig. 2; Collections 42–43).

Description. The evolute shell bears a low rounded keel. The ribs are wiry and widely spaced, reaching a density of up to 20 per half whorl at umbilical diameters greater than 10 mm. The ribs are rectiradiate across the flank and project onto the venter where they abut directly against the keel.

Remarks. This species is more evolute and less densely ribbed than *D. exacta* Buckman, 1905. *D. cf. dumortieri* Thiollière in Dumortier, 1874 is more evolute and slowly expanding, and its ribs are more densely spaced and projecting. *D. varicostata* Géczy, 1967 has less rectiradiate, more projecting ribbing.

Distribution. In Europe, *Dumortieria insignisimilis* is known from the Upper Toarcian of Germany (Ernst 1923), Austria (Fisher 1966), Hungary (Géczy 1967) and possibly Spain (de Villalta and Rosell 1966).

Dumortieria cf. dumortieri (Thiollière in Dumortier, 1874)

Plate 3, figure 16

- cf. 1874 *Ammonites dumortieri* Thiollière in Dumortier; figured by Roman 1938, p. 105, pl. 9, fig. 96; and Arkell *et al.* 1957, p. 262, fig. 296.
 cf. 1892 *Catullocheras dumortieri* (Thiollière); Buckman, p. 277, pl. 39, figs 6–9.
 cf. 1964a *Catullocheras?* sp. indet.; Frebold, p. 16, pl. 7, figs 5–9.
 cf. 1967 *Dumortieria dumortieri* (Thiollière in Dumortier); Géczy, p. 137, pl. 30, fig. 8.
 1968 *Catullocheras cf. C. dumortieri* (Thiollière); Imlay, p. 46, pl. 9, figs 2–4 only.
 cf. 1968 *Dumortieria dumortieri* (Thiollière in Dumortier); Setti, p. 329, pl. 30, fig. 3; pl. 31, fig. 1.
 cf. 1975 *Catullocheras dumortieri* (Thiollière); Guex, p. 115, pl. 7, fig. 1.

Material. Three small and poorly preserved specimens from east-central Oregon (Section 7, locs 5–6). According to Imlay (1968), it is also present in the Bennett area, northern British Columbia (Text-fig. 2; Collections 16–17).

Description. The evolute shell expands slowly and has convex flanks. The ribbing is sharp, dense, rectiradiate, and only weakly projecting onto the venter.

Remarks. These specimens cannot be identified with confidence because of their small size and poor preservation. They are evolute and finely ribbed but not as markedly as, for example, *D. evolutissima* (Prinz, 1904), and we prefer to compare them with *D. dumortieri* which Imlay (1968) has already reported from Oregon on the basis of larger specimens.

Distribution. *Dumortieria dumortieri* is widely distributed in the Upper Toarcian of Europe: in Italy from the Meneghinii Zone (Donovan 1958), in Austria from the Aalensis Subzone of the Levesquei Zone (Fischer 1966), and in France from the Levesquei Subzone of the Pseudoradiosa Zone (Guex 1975; Elmi and Rulleau 1991).

Dumortieria cf. *exacta* Buckman, 1905

Plate 3, figure 15

- cf. 1892 *Dumortieria subundulata* Buckman, pl. 45, figs 6–7 only.
 cf. 1905 *Dumortieria exacta* Buckman, supplement, p. 187.

Material. A single specimen preserved in a siltstone of the Snowshoe Formation, east-central Oregon (Section 7, loc. 5).

Description. A small, moderately evolute specimen ($U = 40$) bearing a low keel. The ribs are sharp and slightly flexuous on early whorls. Ribbing density increases from 21 ribs per half whorl at an umbilical diameter of 6 mm to 29 at 11 mm.

Remarks. As far as the preservation permits comparison, this specimen is similar to a variant of *Dumortieria subundulata* (Buckman, 1892, pl. 45, figs 6–7) that Buckman (1905) elevated to the rank of species and named *D. exacta*. It differs from all other species of *Dumortieria* described from North America by its densely spaced, fine ribbing which is characteristic of the *pseudoradiosa* group of species, as described by Ernst (1923).

Dumortieria raricostata Géczy, 1967

Plate 3, figures 20–21

- 1967 *Dumortieria stefaninii* ? *raricostata* n. subsp. Géczy, p. 142, pl. 31, fig. 8.
 1968 *Dumortieria raricostata* Géczy; Setti, p. 332, pl. 32, fig. 3.
 ?1968 *Catulloceras* cf. *C. dumortieri* (Thiollière); Imlay, p. 46, pl. 9, fig. 5 only.

Material. Three specimens, two of them fragments, preserved in a siltstone of the Snowshoe Formation, east-central Oregon (Section 7, loc. 4).

Description. The shell is evolute, secondarily compressed and bears a low rounded keel. The coarse ribs project onto the venter from the uppermost part of the flank; there are 26 ribs on the outer half whorl. Weak constrictions are evident and the figured specimen bears an 8 mm wide collar that shows faint growth lines and a constricted peristome.

Remarks. The figured specimen represents a mature individual, presumably a macroconch although its microconch is unknown. One specimen of *D. raricostata* is known from the Upper Toarcian of Hungary (Géczy 1967) and two from Italy (Setti 1968), but none show evidence of maturity in spite of reaching larger shell diameters than the Oregon specimen.

Dumortieria? phantasma sp. nov.

Plate 2, figures 3–7; Text-figure 16A–B


Derivation of name. After Phantom Creek in central Graham Island, Queen Charlotte Islands.*Material.* 50 small casts in calcareous concretions and several flattened specimens in shales and siltstones of the Phantom Creek Formation, Queen Charlotte Islands (Section 2, locs 3–5). Also specimens from the Manson River area (Text-fig. 2; Collection 38) and the southern Canadian Rocky Mountains (Collection 45).*Holotype.* GSC 99520 (Pl. 2, figs 3–4) from the lower part of the Phantom Creek Formation (Yakounensis Zone), Yakoun River, Queen Charlotte Islands.*Paratypes.* GSC 99521 (Pl. 2, figs 5–6), GSC 99522 (Pl. 2, fig. 7)

<i>Measurements.</i>	D	UD	U	WH	WW	WWWH
GSC 99520	23.3	8.4	36.1	8.7	6.1	0.70
GSC 99521	18.6	7.5	40.3	6.5	4.6	0.71
GSC 99522	16.9	8.0	47.3	5.2	3.8	0.73

Diagnosis. Compressed, moderately evolute shell; elliptical whorl section; umbilical wall gently sloping; umbilical shoulder gently rounded; flanks gently convex, converging toward sharp venter; ornament generally weak or absent. On outer whorls of some specimens, there are simple, distant, rectiradiate ribs which fade on outer part of flank.*Description.* The holotype, GSC 99520, is a moderately well preserved specimen, septate to 23.1 mm shell diameter with approximately 130° of body chamber ending in an incomplete aperture. The body chamber is slightly crushed. The shell is moderately evolute with a compressed, elliptical whorl section. The flanks are gently convex and converge toward the venter. The venter is sharp but plain. Faint, sparse, rectiradiate ribs are present on the last whorl and fade on the upper part of the flank. The holotype is one of the largest specimens collected, and the sutures on the last part of the phragmocone are approximated. The paratype, GSC 99521, is a moderately well preserved specimen, septate to approximately 5 mm umbilical diameter with approximately 180° of body chamber ending in an incomplete aperture. The body chamber is slightly crushed. The shell has similar features to the holotype. Faint, sparse, rectiradiate ribs are present on the body chamber. The paratype, GSC 99522, is a moderately well preserved specimen, septate to 19.2 mm shell diameter with approximately 100° of body chamber ending in an incomplete aperture. The body chamber is partially crushed.*Remarks.* This form shows similarities to *Atacamiceras* and *Dumortieria*. Its ornament and septal suture are similar to the South American genus *Atacamiceras* described by Hillebrandt (1987). *Atacamiceras glabrum* Hillebrandt, 1987 is smooth except for rare mid-flank ribs. *A. parvicostatum* Hillebrandt, 1987 has smooth inner whorls and is ribbed on the last part of the phragmocone and on the body chamber. Both *Atacamiceras* and the North American form have simple septal sutures

EXPLANATION OF PLATE 4

Figs 1–6, 9–10. *Yakounia silvae* gen. et sp. nov.; Yakounensis Zone, lower part of the Phantom Creek Formation; Queen Charlotte Islands. 1–2, GSC 99525, holotype; GSC Loc. No. C-149652, Section 2, loc. 4. 3–4, GSC 99526, paratype; GSC Loc. No. C-149652, Section 2, loc. 4. 5–6, GSC 99527, paratype; GSC Loc. No. C-149652, Section 2, loc. 4. 9–10, GSC 99528, paratype; GSC Loc. No. C-149652, Section 2, loc. 4.Figs 7–8, 11–14. *Yakounia yakounensis* gen. et sp. nov.; Yakounensis Zone, lower part of the Phantom Creek Formation; Queen Charlotte Islands. 7–8, GSC 99529, paratype; GSC Loc. No. C-149652, Section 2, loc. 4. 11–12, GSC 99530, paratype; GSC Loc. No. C-149652, Section 2, loc. 4. 13–14, GSC 99531, holotype; GSC Loc. No. C-149652, Section 2, loc. 4.Arrows mark start of body chamber. All are $\times 1$.



JAKOBS and SMITH, *Yakounia*

(Text-fig. 16). *Dumortieria? phantasma* has a more compressed shell whereas *Atacamicer* has convex flanks. In addition, *Atacamicer* occurs in the Middle Toarcian of South America whereas the North American form occurs in the latest Toarcian. *Dumortieria? phantasma* is also similar to *D. pusilla* which has smooth internal whorls with distant, simple, rectiradial ribs on the outer whorls (see below). *D. pusilla*, however, has prominent ribbing and achieves it at an earlier stage of development. *D.? phantasma* could be an offshoot of *D. pusilla* via neoteny. The paratype, GSC 99521, displays prominent simple ribs toward the end of the shell, and is similar to the early stage of *D. pusilla*. In addition, the simple septal sutures of *D.? phantasma* could represent the early sutures of *D. pusilla*.

Dumortieria? cf. pusilla Jaworski, 1926

Plate 2, figures 8, 15–16; Plate 3, figure 18

Material. 75 specimens from limestones, mudstones and siltstones of the Snowshoe Formation, east-central Oregon (Section 7, locs 4–5; Section 8, loc. 1), where the species is best represented. Rare and poorly preserved specimens from the Queen Charlotte Islands (Text-fig. 2; Section 3, loc. 97) and the Spatsizi area (Collection 28).

Description. The shell is moderately evolute bearing a weak keel. The ribbing is distant and simple. The ribs are rectiradial on the flanks and project onto the venter where they fade. The inner whorls are smooth.

Discussion. *Dumortieria pusilla* was originally described from Argentina by Jaworski (1926) where it occurs in the South American 'Pleydellia fluitans' Zone, a correlative of the upper Aalensis Subzone of the European Levesquei Zone, according to Hillebrandt (1987). Plaster copies of Jaworski's (1926) type material are figured in Plate 2, figures 9–10 and 13–14. The species is characterized by its smooth inner whorls, a feature that distinguishes it from other species of *Dumortieria* which, at most, have only a small smooth nucleus.

The generic assignment of *Dumortieria pusilla* is based on the simple, distant ribs which characterize many other *Dumortieria* species, such as *D. levesquei* (d'Orbigny, 1844), *D. insignisimilis* and *D. striatulocostata* (Quenstedt, 1885). No other *Dumortieria*, however, possesses smooth internal whorls to such large shell diameters.

Family PHYMATOCERATIDAE Hyatt, 1867
Subfamily PHYMATOCERATINAE Hyatt, 1900


Genus YAKOUNIA gen. nov.

Derivation of name. After the Yakoun River, central Graham Island, Queen Charlotte Islands where Toarcian outcrops are most abundant.

Type species. *Yakounia yakounensis* sp. nov.

EXPLANATION OF PLATE 5

Figs 1–2. *Yakounia freboldi* gen. et sp. nov.; GSC 99532, holotype; Yakounensis Zone, lower part of the Phantom Creek Formation; Queen Charlotte Islands, GSC Loc. No. C-156390, Section 1, loc. 17.
Figs 3–6. *Yakounia yakounensis* gen. et sp. nov.; Yakounensis Zone, lower part of the Phantom Creek Formation; Queen Charlotte Islands. 3–4, GSC 107258, paratype; GSC Loc. No. C-149652, Section 2, loc. 4. 5–6, GSC 107259, paratype; GSC Loc. No. C-149652, Section 2, loc. 4.
Figs 7–8. *Holcophylloceras calypso* (d'Orbigny); GSC 107260; Yakounensis Zone, lower part of the Phantom Creek Formation; Queen Charlotte Islands, GSC Loc. No. C-87221, Section 2, loc. 17.
All are $\times 1$.



JAKOBS and SMITH, *Yakounia* and *Holcophylloceras*

Diagnosis. Moderately evolute shell; ogival whorl section; umbilical wall gently sloping; umbilical shoulder rounded; flanks moderately flat, converging toward a carinate venter with weak ventrolateral shoulders. Venter on inner whorls may be weakly carinate-sulcate. Ornament distinctive, consisting of gently sinuous to approximately rectiradiate ribs that arise in twos or threes from strong umbilical tubercles or prorsiradiate bullae.

Remarks. This form can be compared with several genera of the Phymatoceratinae. According to Donovan (1958), *Brodieia* has sigmoidal ribs joined in groups of two or more to umbilical tubercles. *Brodieia* can be differentiated from *Phymatoceras* and *Haugia* by its non-septate keel. *Yakounia* possesses a septate keel, is more evolute, and the umbilical tubercles are more regular and prominent than in *Brodieia*. *Haugia* has a tall keel, and the ribs and tubercles tend to fade on the outer whorls. In addition, the ribs tend to be more rectiradiate. *Yakounia*'s keel is more subdued, the ornament is prominent on the outer whorls, and the ribs are generally more sinuous and project along the venter. *Phymatoceras* is similar, especially such species as *P. copiapense* and *P. hillebrandti*. In general, however, *Yakounia* has stronger tuberculation, shallower sulci, and a more gently rounded umbilical shoulder. *Yakounia*, especially *Y. pacifica* which has a pronounced angular bend of the ribs low on the flanks, may have evolved into *Ludwigia* or some of the Graphoceratidae.

Distribution. The genus is Late Toarcian in age, occurring with *Hammatoceras*, *Dumortieria*, *Sphaerocoeloceras* and *Pleydellia*. It has been collected from the Queen Charlotte Islands, the Fernie area, southern Alaska, and possibly the Spatsizi area.

Yakounia yakounensis sp. nov.

Plate 4, figures 7-8, 11-14; Plate 5, figures 3-6

- 1976 *Haugia* aff. *H. navis* (Dumortier); Frebold, p. 14, pl. 7, fig. 3.
 1981 *Haugia* cf. *variabilis* (d'Orbigny); Imlay, p. 43, pl. 12, figs 1-2, 5.
 v 1987 ?Grammoceratinae gen. et sp. indet., Hall, p. 1702, pl. 5, figs N-O.

Derivation of name. After the Yakoun River, central Graham Island, Queen Charlotte Islands where Toarcian outcrops are most abundant.

Material. 60 specimens from calcareous concretions and sandstones of the Phantom Creek Formation, Queen Charlotte Islands where the species is best represented (Section 1, loc. 17; Section 2, locs 3-6, 17; Section 3, locs 89, 105; Section 4, locs 2, 4, 6, 14; Section 5, locs 2, 5). Also specimens from the Talkeetna Mountains (Text-fig. 2; Collections 5, 9), the Spatsizi area (Collections 21, 26), and the southern Canadian Rocky Mountains (Collection 45).

Holotype. GSC 99531 (Pl. 4, figs 13-14) from the lower part of the Phantom Creek Formation (middle Yakounensis Zone), Yakoun River, Queen Charlotte Islands.

Paratypes. GSC 99529 (Pl. 4, figs 7-8), GSC 99530 (Pl. 4, figs 11-12), GSC 107258 (Pl. 5, figs 3-4), GSC 107259 (Pl. 5, figs 5-6), GSC 107281-107283

Measurements.	D	UD	U	WH	WW	WWW	PRHW
GSC 99531	64.0	28.0	43.8	19.6	12.3	0.63	10
GSC 99531	50.4	20.0	39.7	17.7	10.7	0.60	9
GSC 99529	65.8	32.4	49.2	17.7	13.0	0.73	11
GSC 99529	54.0	24.7	45.7	17.0	10.5	0.62	10
GSC 107259	40.5	15.8	39.0	15.0	8.5	0.57	10
GSC 107259	33.8	12.0	35.5	13.0	c. 6.6	c. 0.51	10
GSC 107281	54.0	22.4	41.5	17.3	11.6	0.67	10
GSC 107282	68.3	34.0	49.8	20.2	14.7	0.73	12
GSC 107282	53.6	25.0	46.6	16.0	11.2	0.70	10
GSC 107283	110.1	47.5	43.1	36.4	20.7	0.57	13

Diagnosis. Moderately evolute shell; ogival whorl section; umbilical wall gently sloping; umbilical shoulder rounded; flanks flat to gently convex; venter carinate-sulcate on inner whorls becoming carinate on outer whorls; ornament distinctive, consisting of sharp umbilical tubercles from which two to three secondary ribs proceed up the flanks with a gently sinuous trend.

Description. The holotype, GSC 99531, is a well preserved specimen, septate to 55.8 mm shell diameter with approximately 90° of body chamber ending in an incomplete aperture at 64.0 mm shell diameter. The specimen is cracked and a small part of the venter is absent. The shell is moderately evolute with a compressed ogival whorl section. The umbilical wall is gently dipping and the umbilical shoulder is rounded. The flanks are flat to slightly convex and converge toward the carinate venter. The ornament consists of sharp, prominent tubercles at the umbilical shoulder from which two to three secondary ribs proceed up the flank with a gently sinuous trend and terminate at the ventro-lateral shoulder. The tubercles are slightly prorsiradiate and are more prominent on the outer whorls. The paratype, GSC 99530, is a moderately well preserved specimen, septate to 24.6 mm umbilical diameter, with approximately 180° of body chamber ending in an incomplete aperture at approximately 52 mm shell diameter. The specimen is broken on one side. The shell is moderately evolute with a compressed, ogival whorl section. The umbilical wall is gently dipping and the umbilical shoulder is rounded. The flanks are flat to gently rounded. The venter is carinate-sulcate on the inner whorls becoming carinate on the outer whorls. The tubercles at the umbilical shoulder are slightly prorsiradiate and give rise to two or three gently sinuous ribs. The paratypes, GSC 99529 and GSC 107259, are moderately well preserved with parts of their body chambers but incomplete apertures. The body chamber of the latter is slightly crushed. The paratype, GSC 107258, is a moderately well preserved specimen with approximately 230° of body chamber ending in an incomplete aperture. The shell is cracked, slightly crushed and part of the nucleus and venter are absent. The shell is moderately evolute with an ogival whorl section. The whorl section is less compressed than in the previous forms and GSC 107282 has a similar shape. The tubercles at the umbilical shoulder are prominent, sharp and slightly prorsiradiate.

Remarks. This species is distinctive and easily recognized because of its prominent, pointed umbilical tubercles and the gently sinuous ribs. Imlay (1981) compared the southern Alaskan specimens to *Haugia variabilis*, but he noted that they were more evolute and more sparsely ribbed on the outer whorls. He also noted that the holotype of *H. japonica* (Neumayr, 1875) as figured by Kobayashi (1935, pl. 12, figs 3-4) had a much weaker ornament. He compared the southern Alaskan specimens to *Haugia* aff. *japonica* figured by Matsumoto and Ono (1947, pl. 2, fig. 5) but that specimen has much weaker tuberculation, that fades on the outer whorl, and sigmoidal ribbing; it is possibly *Phymatoceras hillebrandti*.

Yakounia yakounensis is distinguished from other species by its prominent umbilical tubercles. *Y. freboldi* sp. nov. has thick, distant ribs with prorsiradiate bullae. *Y. pacifica* sp. nov. has a prominent angular flexure of the ribs, and the ribs and tubercles are subdued. *Y. silvae* sp. nov. is an intermediate form which has characteristics of all three. It possesses tubercles similar to *Y. freboldi*, but the ribbing density is similar to that of *Y. yakounensis*.

Yakounia freboldi sp. nov.

Plate 5, figures 1-2; Text-figure 15c

Derivation of name. After Dr H. Frebold who, for many years, was the Jurassic palaeontologist of the Geological Survey of Canada.

Material. Nine specimens in calcareous concretions and sandstones of the Phantom Creek Formation, Queen Charlotte Islands (Section 1, loc. 17; Section 2, loc. 4; Section 3, loc. 89; Section 4, locs 2, 4, 6; Section 5, locs 2, 5). Also from the southern Canadian Rocky Mountains (Text-fig. 2; Collection 47).

Holotype. GSC 99532 (Pl. 5, figs 1-2) from the lower part of the Phantom Creek Formation (Yakounensis Zone), Yakoun River, Queen Charlotte Islands.

Paratypes. GSC 107284-107287

Measurements.	D	UD	U	WH	WW	WWWH	PRHW
GSC 99532	129.4	57.4	44.4	42.4	30.0	0.71	13
GSC 99532	109.0	48.6	44.6	35.8	25.0	0.70	14
GSC 99532	100.3	44.0	43.9	29.9	20.5	0.69	14
GSC 99532	—	31.3	—	25.5	16.2	0.64	11
GSC 107284	86.0	35.6	41.4	28.3	19.2	0.68	10
GSC 107284	65.8	27.0	41.0	23.1	14.5	0.63	10
GSC 107285	98.0	43.3	44.2	31.1	c. 19.1	c. 0.61	11
GSC 107285	88.4	37.0	41.9	30.7	c. 15	c. 0.49	11
GSC 107285	78.4	32.7	41.7	19.6	11.7	0.60	8
GSC 107286	81.6	30.5	37.4	28.5	19.5	0.68	11
GSC 107287	49.2	16.7	33.9	19.6	11.7	0.60	8
GSC 107287	43.1	14.4	33.4	17.6	11.2	0.64	8

Diagnosis. Moderately evolute shell; ellipsoidal to rectangular whorl section; venter carinate-sulcate on inner whorls becoming carinate on outer whorls; ornament coarse, consisting of prorsiradiate bullae on lower flank from which two to three secondary ribs arise.

Description. The holotype, GSC 99532, is a well preserved specimen with more than 360° of body chamber ending in an incomplete aperture at 129.4 mm shell diameter. Part of the venter is broken away on a small segment of the body chamber. The shell is moderately evolute with a rectangular to ellipsoidal whorl section. The umbilical wall is gently sloping, and the umbilical shoulder is rounded. The flanks are flattish to gently convex. The venter is carinate with weak ventro-lateral shoulders. The ornament is coarse. On the inner whorls, umbilical tubercles give rise to two or three gently sinuous ribs. During ontogeny, the tubercles lengthen into prorsiradiate bullae which give rise to two or three sinuous ribs. The ribs are strongly projecting along the ventro-lateral shoulder and occasionally may reach the venter where subdued swellings are sometimes seen on well preserved parts of the body chamber. The paratype, GSC 107286, is a moderately well preserved specimen wholly septate to a shell diameter of 87.0 mm. The shell is cracked and parts of the phragmocone have broken off. The inner whorls are not exposed. The shell is moderately evolute with a rectangular to ellipsoidal whorl section. The umbilical wall is gently sloping and the umbilical shoulder is rounded. The flanks are flattish, and the venter is carinate with rounded ventro-lateral shoulders. The ornament is coarse with prorsiradiate umbilical bullae, commonly giving rise to two sinuous ribs at approximately one-quarter the flank height. The ribs are strongly projecting along the ventro-lateral shoulder and, where the specimen is well preserved, may reach the venter, where subdued swellings are seen. The other paratypes are not as well preserved but all have similar features to the other types.

Remarks. This species has a coarser ornament than *Yakounia yakounensis*, and the bullae are not as sharp. The inner whorls are similar to *Phymatoceras copiapense* and *Ph. hillebrandti*, which may have been ancestors. *Y. freboldi* may, in turn, be ancestral to *Ludwigia*. The inner whorls have almost straight ribs and sharp tubercles. On the outer whorls, the ribs become more sigmoidal and the tubercles more elongated. The ornament on the outer whorls is similar to that on the inner whorls of species of *Ludwigia* but, in general, that genus has coarser, more sigmoidal ribbing, elongated bullae or primary ribs, and a less sulcate venter.

Yakounia pacifica sp. nov.

Plate 6, figures 1–2

Derivation of name. After the Pacific Ocean, along the north-eastern shore of which the genus is found.

Material. Three specimens in calcareous concretions and sandstones of the Phantom Creek Formation, Queen Charlotte Islands (Section 2, loc. 4).

Holotype. GSC 107261 (Pl. 6, figs 1–2) from the lower part of the Phantom Creek Formation (Yakounensis Zone), Yakoun River, Queen Charlotte Islands.

Paratype. GSC 107288

Measurements.	D	UD	U	WH	WW	WWW	PRHW
GSC 107261	113.4	44.7	39.4	37.5	24.6	0.66	12
GSC 107261	102.7	39.9	38.9	36.0	22.0	0.61	12
GSC 107261	89.5	32.7	36.5	33.6	20.0	0.60	13
GSC 107288	78.4	29.5	37.6	28.3	c. 15.1	c. 0.53	12
GSC 107288	65.8	21.6	32.8	24.4	—	—	11

Diagnosis. Moderately evolute shell; compressed, ogival whorl section; venter carinate-sulcate; ornament prominent, consisting of thick, prorsiradiate primary ribs that split into two or three sinuous secondary ribs low on flanks.

Description. The holotype, GSC 107261, is a moderately well preserved specimen septate to approximately 88 mm shell diameter with approximately 180° of body chamber ending in an incomplete aperture at 113.4 mm shell diameter. The specimen is slightly worn in places. The shell is moderately evolute with an ogival whorl section. The umbilical wall is gently sloping, and the umbilical shoulder is rounded. Flanks are flat, ventrolateral shoulders weak, and the venter carinate. The ornament consists of prorsiradiate primary ribs on the inner whorls that begin high on the umbilical wall. Some ribs may remain single but most bifurcate at approximately one-third the flank height. On the outer whorls, the primary ribs swell into bullae. The paratype, GSC 107288, is a moderately well preserved specimen ending in an incomplete aperture at 78.7 mm shell diameter. The specimen is slightly worn and one side is not exposed. Its morphology is similar to that of the holotype. Septal sutures are present.

Remarks. This species shows some similarity to certain Graphoceratidae such as *Ludwigia* (*Pseudographoceras*) (see Schlegelmilch 1976, pl. 12, figs 5, 7) in terms of whorl shape and ribbing style. However, the Graphoceratidae tend to have a sharp angular flexure of the ribs on the lower flank; this feature is less prominent in similar forms, such as *Phymatoceras hillebrandti*, *Yakounia freboldi*, and *Y. yakounensis*. The Graphoceratidae also tend to be more involute than the Phymatoceratidae although it is possible that they did evolve from them.

Yakounia silvae sp. nov.

Plate 4, figures 1–6, 9–10; Plate 6, figures 3–4

- 1969 *Phlyseogrammoceras* aff. *P. dispansum* (Wunstorff); Frebold *et al.*, p. 20, pl. 1, figs 10–15.
 1987 ?Grammocerotinae gen. et sp. indet., Hall, p. 1702, pl. 5, figs K–L, Q.

Derivation of name. After the thick, old-growth forests that cover(ed) the Queen Charlotte Islands (Latin *silva*, wood, forest).

Material. Over 100 specimens in calcareous concretions and sandstones of the Phantom Creek Formation, Queen Charlotte Islands (Section 1, locs 4–5, 7–8, 17; Section 2, locs 1–6, 8, 10–11, 17; Section 3, locs 89–92, 105; Section 4, locs 1–2, 4, 6–8, 10, 14; Section 5, locs 2–3, 5; Section 6, locs 1–2). Also specimens from the Manson River area (Text-fig. 2; Collection 38), in the Manning Park area (Text-fig. 2; Collection 44), and in the southern Canadian Rocky Mountains (Collection 45).

Holotype. GSC 99525 (Pl. 4, figs 1–2) from the lower part of the Phantom Creek Formation (Yakounensis Zone), Yakoun River, Queen Charlotte Islands.

Paratypes. GSC 99526 (Pl. 4, figs 3–4), GSC 99527 (Pl. 4, figs 5–6), GSC 99528 (Pl. 4, figs 9–10), GSC 107262 (Pl. 6, figs 3–4), GSC 107289–107291


Measurements.	D	UD	U	WH	WW	WWWH	PRHW
GSC 99525	82.0	31.0	37.8	29.2	c. 18	c. 0.62	11
GSC 99525	66.9	23.7	35.4	23.7	14.9	0.63	11
GSC 107289	79.0	32.8	41.5	27.0	16.0	0.59	10
GSC 107289	86.1	38.1	44.3	27.2	16.7	0.61	11
GSC 107290	102.9	39.0	37.9	37.0	22.0	0.59	11
GSC 107291	60.6	25.0	41.3	19.5	12.8	0.66	12
GSC 107291	53.0	21.3	40.2	18.6	11.4	0.61	12

Diagnosis. Moderately evolute planulate; ogival whorl section; venter carinate-sulcate; ornament prominent, consisting of umbilical tubercles or prorsiradiate bullae from which two or three secondary ribs proceed up the flanks with a gently sinuous to rectiradiate trend.

Description. The holotype, GSC 99525, is a moderately well preserved specimen wholly septate to a shell diameter of 83.0 mm. Part of one side of the outer whorl is crushed. The shell is moderately evolute with a compressed ogival whorl section. The umbilical wall is gently sloping and the umbilical shoulder is rounded. The flanks are moderately flat to gently convex. The venter is carinate with weak ventro-lateral shoulders. On the inner whorls, prorsiradiate primaries begin high on the umbilical wall and split into two or three primaries at approximately one-third the flank height. On the outer whorls, prorsiradiate umbilical tubercles develop and on the outer whorl are bullate. The tubercles/bullae give rise to two or three gently sinuous secondary ribs. The paratype, GSC 107262, is a moderately well preserved specimen with more than 180° of body chamber ending in an incomplete aperture at 96.9 mm shell diameter. The specimen, especially the body chamber, is partially crushed. The shell is moderately evolute. The umbilical wall is gently sloping with a rounded umbilical shoulder. The flanks are flattish and the venter is carinate. On the inner whorls, umbilical tubercles give rise to two or three secondary ribs. The tubercles become slightly prorsiradiate on the outer whorl. The secondary ribs are almost rectiradiate to gently sinuous. The paratype, GSC 99528, is a moderately well preserved specimen septate to 13.3 mm umbilical diameter with approximately 200° of body chamber ending in an incomplete aperture at 42.8 mm shell diameter. The body chamber is partially crushed. The shell form is similar to those described above. On the inner whorls, gently prorsiradiate primary ribs give rise to two or three secondary ribs at approximately one-third the flank height. Umbilical tubercles develop on the last part of the phragmocone and on the body chamber. The secondary ribs are rectiradiate to gently sinuous. The paratype, GSC 99527, is a moderately well preserved specimen with part of the body chamber ending in an incomplete aperture at 35.3 mm shell diameter. Part of the phragmocone venter is absent, and the body chamber near the aperture is partially crushed. The tubercles begin on the last whorl, and the secondary ribs are almost rectiradiate. The paratype, GSC 99526, is a moderately well preserved specimen, septate to 33.3 mm shell diameter with approximately 160° of body chamber ending in an incomplete aperture at 43.5 mm shell diameter. The body chamber is partially crushed. Tubercles begin on the last whorl, and give rise to two or three gently sinuous secondary ribs.

EXPLANATION OF PLATE 6

- Figs 1–2. *Yakounia pacifica* gen. et sp. nov; GSC 107261, holotype; Yakounensis Zone, lower part of the Phantom Creek Formation; Queen Charlotte Islands, GSC Loc. No. C-149652, Section 2, loc. 4.
 Figs 3–4. *Yakounia silvae* gen. et sp. nov; GSC 107262, paratype; Yakounensis Zone, lower part of the Phantom Creek Formation; Queen Charlotte Islands, GSC Loc. No. C-149652, Section 2, loc. 4.
 Figs 5–6. *Pseudolioceras compactile* (Simpson); GSC 107263; Yakounensis Zone, lower part of the Phantom Creek Formation; Queen Charlotte Islands, GSC Loc. No. C-157742, Section 2, loc. 95.
 Figs 7–10. *Sphaeroceoloceras brochiiforme* Jaworski; Yakounensis Zone, lower part of the Phantom Creek Formation; Queen Charlotte Islands. 7–8, GSC 95584; GSC Loc. No. C-87118, Section 3, loc. 95. 9–10, GSC 99485; GSC Loc. No. C-149652, Section 2, loc. 4.
 Arrow marks start of body chamber. All are $\times 1$.



JAKOBS and SMITH, *Yakounia*, *Pseudolioceras* and *Sphaerocoeloceras*

Remarks. This species has a weaker ornament than *Y. frebaldi*, although the prorsiradiate bullae are similar. It lacks the sharp, round tubercles of *Y. yakounensis* and has more prominent tubercles than *Y. pacifica*. In general, the specimens belonging to *Y. silvae* are alike except that certain features may arise or fade at different points during ontogeny, and ribs and bullae may trend in slightly different directions.

Subfamily HARPOCERATINAE Neumayr, 1875

Genus PSEUDOLIOCERAS Buckman, 1889

[= *Praehaploceras* Monestier, 1931; *Pseudowalkeriaceras* Maubeuge, 1949; *Tugurites* Kalacheva and Sey, 1970; ?*Buckmanites* Guex, 1973]

Type species. *Ammonites compactilis* Simpson, 1855 (subsequent designation by Buckman 1889)

Diagnosis. Moderately involute shell with discoidal and compressed triangular whorl section; carinate venter with sharp, hollow keel bordered by narrow, smooth areas; umbilical wall vertical; umbilicus small and deep; ribs sub-falciform, only clearly visible on outer part of whorl where they are broad and rounded; on inner part of whorl, ribs either faint or absent.

Distribution. *Pseudolioceras* appears to have Boreal affinities, being common in Arctic Canada, Siberia, Japan and Great Britain.

Pseudolioceras compactile (Simpson, 1855)

Plate 6, figures 5–6

- *1855 *Ammonites compactilis* Simpson, p. 75.
- 1911 *Pseudolioceras compactile* (Simpson); Buckman, pp. 41b, 41c, pl. 41A, figs 1–2.
- 1972 *Pseudolioceras compactile* (Simpson); Guex, pl. 5, fig. 13.
- 1974 *Pseudolioceras compactile* (Simpson); Dagens, pl. 18, fig. 1.
- 1976 *Pseudolioceras compactile* (Simpson); Schlegelmilch, p. 89, pl. 47, fig. 4.

Material. 11 small specimens in sandstones and calcareous concretions of the Phantom Creek Formation, Queen Charlotte Islands (Section 3, loc. 95; Section 4, loc. 12).

<i>Measurements.</i>	D	UD	U	WH	WW	WWWH	PRHW
GSC 107263	22.8	3.7	16.2	13.2	6.9	0.52	—
GSC 107292	18.9	3.1	16.4	10.5	—	—	—

Description. The involute shell has a triangular whorl section. The umbilical shoulder is rounded, and the umbilical wall steep. The flanks are gently convex. The venter is almost knife-sharp; the flanks converge toward the venter with weak ventro-lateral shoulders. The ribs are gently falcate, projecting along the venter. On the lower flanks, they are faint, becoming more prominent on the upper flank. They are much fainter on the early whorls, appearing almost like growth lines.

Remarks. The degree of involution, the discoidal nature of the shell, and the knife-edge venter of the Queen Charlotte Islands specimens are typical of *P. compactile*. *P. lythense* (Young and Bird, 1828) has coarser ribbing that is more prominent on the lower flanks. Some specimens are similar to *Leioceras opalinum* (Reinecke, 1818) with fine ribbing, and a similar whorl shape and volution. These could be juvenile features of *Pseudolioceras*, as they appear on specimens of small size.

Distribution. *Pseudolioceras compactile* has been collected from the Upper Toarcian of north-west Europe, and is common in the Arctic regions of North America and Siberia.

Subfamily HAMMATOCERATINAE Buckman, 1887

Genus HAMMATOCERAS Hyatt, 1867

[= *Ammatoceras* Hyatt, 1867; *Pachammatoceras* Buckman, 1921]*Type species.* *Ammonites insignis* Zieten, 1831 (subsequent designation by Buckman 1887)*Diagnosis.* Moderately evolute shell with sub-triangular to ogival whorl section; umbilical wall becomes vertical and commonly undercut on outer whorls; primary ribs short, with long secondary ribs arising near umbilical shoulder, commonly at tubercles; venter carinate; prorsiradiate ribs approach it at slight angle; ribs fade on outer whorls.*Distribution.* With the exception of the Arctic regions, *Hammatoceras* is a cosmopolitan genus found in the Upper Toarcian and Lower Aalenian.*Hammatoceras insigne* (Zieten, 1831)

Plate 7, figure 5

- 1831 *Ammonites insignis* Zieten, pl. 15, fig. 2.
 1874 *Ammonites insignis* (Schubler); Dumortier, pl. 17, figs 1-3 (*non* 4-5).

Material. Only three specimens known, collected from calcareous concretions and sandstones of the Phantom Creek Formation, Queen Charlotte Islands (Section 2, loc. 4, Section 3, locs 71, 95).*Description.* The shell is involute with a triangular whorl section. Flanks are convex to the carinate venter. On the outer whorls, the keel is bordered by smooth bands. The ornament is prominent with sharp umbilical tubercles from which two or three rectiradiate secondary ribs arise which approach the venter at an angle of approximately 90°.*Remarks.* This species can be distinguished from *H. speciosum* by the latter's more compressed whorl section, and ribs that approach the venter with a slightly prorsiradiate trend.*Distribution.* *Hammatoceras insigne* is well known from the Upper Toarcian of Europe (Guex 1975; Schlegelmilch 1976; Elmi and Rulleau 1991).*Hammatoceras speciosum* (Janensch, 1902)

Plate 7, figures 1-4, 6

- 1874 *Ammonites insignis* (Schubler); Dumortier, pl. 18.
 1885 *Ammonites insignis variabilis* Quenstedt, pl. 50, fig. 1.
 *1902 *Ammonites speciosum* Janensch, pl. 10, fig. 1.
 1974 *Hammatoceras speciosum* (Janensch); Elmi *et al.*, pl. 5, fig. 1.
 1975 *Hammatoceras speciosum* (Janensch); Guex, p. 111, pl. 9, fig. 8; pl. 10, figs 1, 3.
 1976 *Hammatoceras speciosum* (Janensch); Schlegelmilch, p. 91, pl. 49, fig. 2.
 1991 *Hammatoceras speciosum* (Janensch); Tipper *et al.*, pl. 7, fig. 1.
 1992 *Hammatoceras speciosum* (Janensch); Hillebrandt and Smith, pl. 4, fig. 1.

Material. 20 specimens from calcareous concretions and sandstone of the Phantom Creek Formation, Queen Charlotte Islands (Section 1, loc. 3; Section 2, loc. 4; Section 3, locs 90, 99-100, 105; Section 4, loc. 2; Section 5, locs 1-2, 5). Two poorly preserved specimens from the Snowshoe Formation, east-central Oregon (Text-fig. 2; Section 7, locs 4-5).

Measurements.	D	UD	U	WH	WW	WWW	PRHW
GSC 95583	130.8	55.0	42.0	43.3	34.8	0.80	18
GSC 95583	119.2	50.6	42.4	39.5	32.0	0.81	17
GSC 107264	29.3	9.6	32.8	12.4	11.4	0.92	9
GSC 107264	23.6	7.0	29.7	10.0	9.8	0.98	8
GSC 107293	—	—	—	23.0	17.0	0.74	—
GSC 107294	133.1	—	—	45.1	38.5	0.85	—
GSC 107295	c. 135	55.0	40.7	45.5	—	—	—
GSC 107295	92.4	33.9	36.7	33.3	—	—	12

Description. The shell is moderately evolute with an ogival to triangular whorl section. The umbilical wall is steep, becoming vertical and undercut on the outer whorls. The umbilical shoulder is rounded but abrupt. The flanks are gently convex, converging toward the carinate venter. The ornament is coarse, consisting of strong umbilical tubercles from which two or three secondary ribs arise. The primary ribs are rursiradial on the umbilical wall. The secondary ribs are gently curved and approach, but do not reach, the venter at an angle of approximately 70°.

Remarks. The whorls are much taller and narrower than those of *Hammatoceras insigne*, and the ribbing is finer and denser. In addition, the ribs of *H. insigne* approach the venter at an angle of almost 90°. *H. bonarelli* Parisch and Viale, 1906 is more evolute, and the whorls are not as high as those of *H. speciosum*.

Distribution. *Hammatoceras speciosum* is common in Europe and the Mediterranean region: in Algeria from the Insigne Zone (Elmi *et al.* 1974), in southern Germany from the Levesquei Zone (Schlegelmilch 1976), and in France from the Insigne Zone (Guex 1975; Elmi and Rulleau 1991).

Hammatoceras sp. nov.

Plate 7, figures 7–8

Material. Three incomplete specimens in sandstones and calcareous concretions of the Phantom Creek Formation, Queen Charlotte Islands (Section 4, loc. 6; Section 5, loc. 2)

Measurements.	D	UD	U	WH	WW	WWW	PRHW
GSC 107296	—	—	—	12.7	13.3	1.05	—
GSC 107297	—	—	—	9.5	14.0	1.47	—

Description. The shell is moderately evolute with a depressed, almost wide oval whorl section. The flanks are flat, becoming convex on the outer whorls. The venter is carinate with a subdued keel. The ornament is coarse and sparse. Umbilical tubercles give rise to two or three secondary ribs which are rectiradial on the flank, but become prorsiradial and approach the venter at an angle of approximately 70°.

EXPLANATION OF PLATE 7


Figs 1–4. *Hammatoceras speciosum* Janensch; Yakounensis Zone, lower part of the Phantom Creek Formation; Queen Charlotte Islands. 1, 4, GSC 95583; GSC Loc. No. C-87215, Section 2, loc. 4. 2–3, GSC 107264, GSC Loc. No. C-87216, Section 2, loc. 4.

Fig. 5. *Hammatoceras insigne* (Zieten); GSC 107265; Hillebrandti Zone, upper part of the Whiteaves Formation; Queen Charlotte Islands, GSC Loc. No. C-159396, Section 3, loc. 71.

Fig. 6. *Hammatoceras speciosum* Janensch; UBC 017; Yakounensis Zone, Warm Springs Member of the Snowshoe Formation; Izee area, eastern Oregon, UBC Loc. No. F4-4-E, Section 7, loc. 5.

Figs 7–8. *Hammatoceras* sp. nov.; GSC 107266; Yakounensis Zone, lower part of the Phantom Creek Formation; Queen Charlotte Islands, GSC Loc. No. C-158077, Section 5, loc. 2.

All are × 1.



JAKOBS and SMITH, *Hammatoceras*

Remarks. This species is similar to *H. insigne* but differs in possessing much coarser and more distant ribbing, and in that the ribs approach the venter at a more acute angle.

Genus *SPHAEROCOELOCERAS* Jaworski, 1926

Type species. *Sphaerocoeloceras brochiiforme* Jaworski, 1926

Diagnosis. Small involute shell with wide ellipsoidal to rounded whorl section; venter plain, but sometimes weakly carinate; ribbing blunt, sparse, bifurcating low on flank; no tubercles present.

Remarks. This genus has been known previously only from South America. It was placed initially within the Dactylioceratidae by Jaworski (1926) because of the similarity of its suture line to that of *Coeloceras crassum* Young and Bird, 1828. Arkell *et al.* (1957) retained this classification. Donovan *et al.* (1981) placed it within the Hammatoceratinae. It is similar to small specimens of *Hammatoceras* in appearance except for the missing keel. The genus is of Late Toarcian age in South and North America.

Sphaerocoeloceras brochiiforme Jaworski, 1926

Plate 6, figures 7–10

- *1926 *Sphaerocoeloceras brochiiforme* Jaworski, p. 259, pl. 1, fig. 10; pl. 3, fig. 10; pl. 4, fig. 23.
- v 1987 *Sphaerocoeloceras brochiiforme* Jaworski; Hillebrandt, pl. 13, fig. 9a–c; pl. 14, fig. 10a–b.
- v 1987 ?Hammatoceratidae gen. et sp. indet. Hall, p. 1702, pl. 5, figs R–V.
- v 1991 *Sphaerocoeloceras brochiiforme* Jaworski; Tipper *et al.*, pl. 7, fig. 2.
- v 1992 *Sphaerocoeloceras brochiiforme* Jaworski; Hillebrandt and Smith, pl. 4, fig. 2.
- v 1994 *Sphaerocoeloceras brochiiforme* Jaworski; Jakobs *et al.*, pl. 5, figs 7–8.

Material. 40 specimens in calcareous concretions and sandstones of the Phantom Creek Formation, Queen Charlotte Islands (Section 1, loc. 17; Section 2, locs 3–4, 17; Section 3, locs 95, 97–98; Section 4, locs 2, 6, 10; Section 5, loc. 5). Poorly preserved specimens from the southern Canadian Rocky Mountains (Text-fig. 2; Collection 45).

Measurements.	D	UD	U	WH	WW	WWWH	PRHW
GSC 107298	20.4	7.9	38.7	5.0	9.5	1.90	14
GSC 107299	17.8	6.7	37.6	4.6	8.6	1.87	15
GSC 107300	16.2	5.0	30.9	5.5	9.0	1.64	14
GSC 107301	24.7	9.7	39.3	7.4	9.9	1.34	16
GSC 107302	17.0	6.2	36.5	5.7	10.7	1.88	13
GSC 107303	19.0	7.0	36.8	5.0	8.2	1.64	15

Description. The shell is moderately evolute with a depressed, rounded whorl section. The umbilical wall is gently sloping and the umbilical shoulder is weak. The flanks are gently convex, converging towards the plain venter. No keel is evident but a faint ridge may sometimes be present. The ornament is generally faint. Ribs are blunt and begin near the umbilical shoulder, prorsiradiate to the upper flank, then rursiradiate to the venter. The ribs approach the venter with an angle of slightly less than 180°, and their ventral ends may be separated by a smooth area. They commonly bifurcate at the top of the flank, and single intercalatory ribs may also appear. The venter is similar to that of *Hammatoceras* except for the absence of a keel. Ribbing strength and density vary slightly between the specimens.

Remarks. Slight variations in ribbing density and strength are not considered sufficiently important to warrant splitting the group into several species.

Distribution. *Sphaerocoeloceras brochiiforme* has been collected in South America, from the *Phlyseogrammoceras* (?) *tenuicostatum*, '*Pleydellia lotharingica*' and '*Pleydellia fluitans*' zones of Hillebrandt (1987).

CONCLUSIONS

The latest Toarcian ammonite fauna of North America contains endemic Athabascan (*sensu* Taylor *et al.* 1984) and eastern Pacific taxa, as well as pandemic taxa. Eastern Pacific taxa such as *Sphaerocoeloceras brochiiforme* and *Dumortieria pusilla* provide correlative ties with South American faunas described by Hillebrandt (1981, 1987). Pandemic taxa, such as *Pleydellia aalensis*, *Hammatoceras speciosum*, *Dumortieria levesquei*, and *D. insignisimilis*, allow correlation with European successions. Athabascan forms are the most widespread and prolific taxa in western North America; these include *Yakounia yakounensis*, *Y. frebaldi*, *Y. silvae*, and *Pleydellia maudensis*. A global sea-level fall during the Late Toarcian (Hallam 1988; Haq *et al.* 1988) may have encouraged endemism by restricting migration between the eastern Pacific and western Tethys.

Acknowledgements. Most of this study was part of the doctoral research of GKJ, supported by a NSERC research grant to PLS and by the Geological Survey of Canada. We thank H. W. Tipper for making available old collections and sharing his extensive knowledge of the Canadian Jurassic, Beth Carter for identifying Radiolaria, and Dave Taylor for identifying Middle Jurassic ammonites. Photographs were taken by K. Gordanier-Smith.

REFERENCES

- ANDERSON, R. G. and THORKELSON, D. J. 1990. Mesozoic stratigraphy and setting for some mineral deposits in Iskut River map area, northwestern British Columbia. In *Current Research, Part E, Geological Survey of Canada Paper*, **90-1F**, 131–139.
- ARKELL, W. J. 1950. A classification of the Jurassic ammonites. *Journal of Paleontology*, **24**, 354–364.
- KUMMEL, B. and WRIGHT, C. W. 1957. Mesozoic ammonoidea. L80–L490. In MOORE, R. C. (ed.). *Treatise on invertebrate paleontology. Part L. Mollusca 4*. Geological Society of America and University of Kansas Press, Boulder, Colorado and Lawrence, Kansas, 490 pp.
- ARTHUR, A., SMITH, P. L., MONGER, J. W. H. and TIPPER, H. W. 1993. Mesozoic stratigraphy and Jurassic paleontology west of Harrison Lake, southwestern British Columbia. *Bulletin of the Geological Survey of Canada*, **441**, 63 pp., 5 pls.
- BAYLE, E. 1878. Fossiles principaux des terraines. *Service de la Carte géologique détaillée. Explication de la Carte géologique détaillée*, **4**, Atlas, part 1, 158 pls.
- BENECKE, E. W. 1865. Über Trias und Jura in den Südalpen. *Geognostische-Paläontologische Beiträge*, **1**, 1–204, pls 1–11.
- BRANCO, W. 1879. Der Untere Dogger Deutsche-Lothringens. *Abhandlungen zur Geologischen Spezialkarte von Elsass-Lothringen*, **2(2)**, 1–160, pls 1–10.
- BRAUNS, D. 1865. Die Stratigraphie und Paläontologie des südöstlichen Theiles der Hilmulde. *Palaeontographica*, **13**, 75–146.
- BUCKMAN, S. S. 1887–1907. A monograph on the Inferior Oolite ammonites of the British Islands. *Monograph of the Palaeontographical Society London*, Part 1, 1887, **40(192)**, 1–24, pls 1–6; Part 2, 1888, **41(197)**, 25–56, pls 7–14; Part 3, 1889, **42(201)**, 57–144, pls 15–23A; Part 4, 1890, **43(206)**, 145–224, pls 24–36; Part 5, 1891, **44(210)**, 225–256, pls 37–44; Part 6, 1892, **45(214)**, 257–312, pls 45–56; Part 7, 1892, **46(220)**, 313–344, pls 57–76; Part 8, 1893, **47(224)**, 345–376, pls 77–92; Part 9, 1894, **48(228)**, 377–456, pls 93–103; Part 10, 1898, **52(245)**, 1–32, pls 1–4; Part 11, 1899, **53(251)**, 33–64, pls 5–14; Part 12, 1904, **58(275)**, 65–168, pls 15–19; Part 13, 1905, **59(281)**, 169–208, pls 20–24; Part 14, 1907, **61(292)**, 209–262.
- 1909–1930. *Yorkshire Type Ammonites* (2 vols) and *Type Ammonites* (5 vols). William Wesley and Son, London, 790 pls.
- CAMERON, B. E. B. and TIPPER, H. W. 1985. Jurassic stratigraphy of the Queen Charlotte Islands, British Columbia. *Bulletin of the Geological Survey of Canada*, **365**, 49 pp.
- CARTER, E. S. and JAKOBS, G. K. 1990. New Aalenian Radiolaria from the Queen Charlotte Islands, British Columbia: implications for biostratigraphic correlation. In *Current Research, Part A, Geological Survey of Canada Paper*, **91-1A**, 337–351.
- DAGYS, A. A. 1974. Toarskie ammonity (Hildoceratidae) severa Sibiri. *Trudy Instituta Geologii i Geofiziki, Novosibirsk*, **99**, 108 pp., 19 pls. [In Russian].

- DENCKMANN, A. 1887. Über geognostischen Verhältnisse der Umgegend von Dörnten nordlich Goslar, mit besonderer Berücksichtigung der Fauna des oberen Lias. *Atlas zu den Abhandlungen zur geologischen Spezialkarte von Preussen und den Thüringischen Staaten*, 8(2), Berlin, 108 pp., 10 pls.
- DICKINSON, W. R. and VIGRASS, L. W. 1964. Pre-Cenozoic history of the Suplee-Izee district, Oregon. *Bulletin of the State of Oregon, Department of Geology and Mineral Industries*, 58, 109 pp.
- DONOVAN, D. T. 1958. The ammonite zones of the Toarcian (Ammonitico Rosso Facies) of southern Switzerland and Italy. *Eclogae geologicae Helvetiae*, 51, 33–60.
- CALLOMON, J. H. and HOWARTH, M. K. 1981. Classification of the Jurassic Ammonitina. 101–155. In HOUSE, M. R. and SENIOR, J. R. (eds). *The Ammonoidea*. Systematics Association London, Special Volume, 18.
- DUMORTIER, E. 1984. *Études paléontologiques sur les dépôts jurassiques du Bassin du Rhône, 4, Lias Supérieur*. Paris, 335 pp., 65 pls.
- ELMI, S. and RULLEAU, L. 1987. Le genre *Pseudolillia* (Ammonitina, Grammocerotinae) dans le Toarcien Supérieur (France, Espagne, Portugal, Maroc). *Atti del secondo convegno internazionale – Fossili, Evoluzione, Ambiente*, 291–315, pls 1–7.
- 1991. Le Toarcien des Carrières Lafarge (Bas-Beaujolais, France): cadre biostratigraphique de référence pour la région Lyonnaise. *Geobios*, 24, 315–331, pls 1–5.
- ATROPS, F. and MANGOLD, C. 1974. Les zones d'ammonites du Domérien–Callovien de l'Algérie Occidentale – Première partie: Domérien–Toarcien. *Documents des Laboratoires de Géologie de la Faculté des Sciences de Lyon*, 61, 1–83, pls 1–6.
- ERNST, W. 1923. Zur stratigraphie und fauna des Lias zeta in nordwestlichen Deutschland. *Palaeontographica*, 65, 1–95, pls 1–6.
- FISCHER, R. 1966. Die Dactyloceratidae (Ammonoidea) der Kammerker (Nord-tirol) und die Zonengliederung des alpinen Toarcien. *Bayerische Akademie der Wissenschaften, Mathematisch-Naturwissenschaftliche Klasse, Abhandlung*, 126, 83 pp., 6 pls.
- FREBOLD, H. 1957. The Jurassic Fernie Group in the Canadian Rocky Mountains and foothills. *Memoir of the Geological Survey of Canada*, 287, 197 pp., 44 pls.
- 1964a. Lower Jurassic and Bajocian ammonoid faunas of northwestern British Columbia and southern Yukon. *Bulletin of the Geological Survey of Canada*, 116, 31 pp., 8 pls.
- 1964b. Illustrations of Canadian fossils. Jurassic of Arctic and Western Canada. *Geological Survey of Canada Paper*, 63–4, 107 pp., 51 pls.
- 1969. Subdivision and facies of Lower Jurassic rocks in the southern Canadian Rocky Mountains and foothills. *Proceedings of the Geological Association of Canada*, 20, 76–89, 2 pls.
- 1976. The Toarcian and Lower Middle Bajocian beds and ammonites in the Fernie Group of southeastern British Columbia and parts of Alberta. *Geological Survey of Canada Paper*, 75–39, 20 pp., 9 pls.
- MOUNTJOY, E. W. and TEMPELMAN-KLUIT, D. J. 1967. New occurrences of Jurassic rocks and fossils in central and northern Yukon Territory. *Geological Survey of Canada Paper*, 67–12, 35 pp., 3 pls.
- and TIPPER, H. W. 1970. Status of the Jurassic in the Canadian Cordillera of British Columbia, Alberta and Southern Yukon. *Canadian Journal of Earth Sciences*, 7, 1–21.
- and COATES, J. A. 1969. Toarcian and Bajocian rocks and guide ammonites from southwestern British Columbia. *Geological Survey of Canada Paper*, 67–10, 55 pp., 6 pls.
- FRECH, F. 1902. Über devonische Ammoneen. *Beiträge zur Paläontologie und Geologie von Österreich-Ungarn und Orients*, 14, 27–112, pls 2–5.
- GÉCZY, B. 1967. Ammonoidés jurassiques de Csernye, Montagne Bakony, Hongrie. Part II (excl. Hammatoceratidae). *Geologica Hungarica, Series Palaeontologica*, 35, 413 pp., 65 pls.
- GEMMELLARO, G. 1886. Monografia sui fossili del Lias superiore delle provincie di Palermo. *Giornale di Scienze naturali ed economiche Palermo*, 17, 188–197.
- GOY, A. and MARTÍNEZ, G. 1990. Biozonación del Toarciense en el área de La Almunia de Doña Godina-Ricla (Sector Central de la Cordillera Ibérica). *Cuadernos de Geología Ibérica*, 14, 11–53, pls 1–4.
- GUEX, J. 1972. Répartition biostratigraphique des ammonites du Toarcien moyen de la bordure sud des Causses (France) et révision des ammonites décrites et figurées par Monestier (1931). *Eclogae geologicae Helvetiae*, 65, 611–645, 13 pls.
- 1973. Observations sur la répartition biostratigraphique des ammonites du Toarcien supérieur de l'Aveyron (France). *Bulletin des Laboratoires de Géologie, Minéralogie, Géophysique et du Musée Géologique de l'Université de Lausanne*, 207, 7–14, pls 1–2.
- 1975. Description biostratigraphique du Toarcien supérieur de la bordure sud des Causses (France). *Eclogae geologicae Helvetiae*, 68, 97–129, pls 1–12.

- HALL, R. L. 1984. Lithostratigraphy and biostratigraphy of the Fernie Formation (Jurassic) in the Southern Canadian Rocky Mountains. In STOTT, D. F. and GLASS, D. J. (eds). The Mesozoic of Middle North America. *Memoir of the Canadian Society of Petroleum Geologists*, **9**, 233–247.
- 1987. New Lower Jurassic ammonite fauna from the Fernie Formation, southern Canadian Rocky Mountains. *Canadian Journal of Earth Sciences*, **24**, 1688–1704, 5 pls.
- HALLAM, A. 1988. A re-evaluation of Jurassic eustasy in light of new data and the revised Exxon curve. *Sea-level changes: an integrated approach*, *Society of Economic Paleontologists and Mineralogists, Special Publication*, **42**, 261–273.
- HAQ, B. U., HARDENBOL, J. and VAIL, P. R. 1988. Mesozoic and Cenozoic chronostratigraphy and cycles of sea-level change. *Sea-level changes: an integrated approach*, *Society of Economic Palaeontologists and Mineralogists, Special Publication*, **42**, 71–108.
- HAUG, E. 1885. Beiträge zu einer Monographie der Ammonitagattung *Harpoceras*. *Neues Jahrbuch für Mineralogie, Geologie und Paläontologie, Beilage-Band*, **3**, 585–722, pls 11–12.
- HILLEBRANDT, A. von 1981. Kontinentalverschiebung und die paläozoogeographischen Beziehungen des südamerikanischen Lias. *Geologische Rundschau*, **70**, 570–582.
- 1987. Liassic ammonite zones of South America and correlations with other provinces with descriptions of new genera and species of ammonites. *Biostratigrafía de los sistemas regionales del Jurásico y Cretácico en América del Sur, Tomo I: Jurásico anterior a los movimientos málmicos*, **1**, 111–157, 14 pls.
- and SMITH, P. L. 1992. Ammonite zones of the circum-Pacific region: Lower Jurassic. Chapter 12, 247–253. In WESTERMANN, G. E. G. (ed.). The Jurassic of the Circum-Pacific. *International Geological Correlation Programme Project 171: Jurassic of the Circum-Pacific*. Cambridge University Press, 676 pp.
- HYATT, A. 1867. The fossil Cephalopoda of the Museum of Comparative Zoology. *Bulletin of the Museum of Comparative Zoology*, **5**, 71–102.
- 1889. Genesis of the Arietidae. *Smithsonian Contributions to Knowledge, Washington*, **673**, xi + 238 pp., 14 pls.
- 1900. Cephalopoda. 502–592, figs 1049–1235. In ZITTEL, K. A., *Textbook of Palaeontology*, 1st English edition, translated by Eastman, C. R.
- IMLAY, R. W. 1968. Lower Jurassic (Pliensbachian and Toarcian) ammonites from eastern Oregon and California. *United States Geological Survey Professional Paper*, **593-C**, 51 pp., 9 pls.
- 1973. Middle Jurassic (Bajocian) ammonites from eastern Oregon. *United States Geological Survey Professional Paper*, **756**, 100 pp., 48 pls.
- 1981. Early Jurassic ammonites from Alaska. *United States Geological Survey Professional Paper*, **1148**, 49 pp., 12 pls.
- IRVING, E. and WYNNE, P. J. 1991. Paleomagnetism: review and tectonic implications. Chapter 3, 61–86. In GABRIELSE, H. and YORATH, C. J. (eds). Geology of the Cordilleran Orogen in Canada. *Geological Survey of Canada, Geology of Canada, no. 4 (also Geological Society of America, The Geology of North America, v. G-2)*, 844 pp.
- and YOLE, R. W. 1987. Tectonic rotations and translations in western Canada; new evidence from Jurassic rocks of Vancouver Island. *Geophysical Journal of the Royal Astronomical Society*, **91**, 1025–1048.
- MONGER, J. W. H. and YOLE, R. W. 1980. New paleomagnetic evidence for displaced terranes in British Columbia. 441–456. In STRANGWAY, D. W. (ed.). The continental crust and its mineral deposits. *Geological Association of Canada, Special Paper*, **20**, 804 pp.
- JAKOBS, G. K. 1990. A discussion of the Phantom Creek Formation of the Maude Group. Queen Charlotte Islands, British Columbia. In *Current Research, Part F, Geological Survey of Canada Paper*, **90-1F**, 57–60.
- 1992. Toarcian (Lower Jurassic) ammonite biostratigraphy and ammonite fauna of North America. Unpublished Ph.D. thesis, University of British Columbia, 682 pp., 23 pls.
- SMITH, P. L. and TIPPER, H. W. 1994. An ammonite zonation for the Toarcian (Lower Jurassic) of the North American Cordillera. *Canadian Journal of Earth Sciences*, **31**, 919–942, pls 1–5.
- — — 1995 (dated 1994). Towards an ammonite zonation for the Toarcian of North America. Third International Symposium on Jurassic Stratigraphy, Poitiers, France. *Geobios, Mémoire Spécial*, **17**, 317–325.
- JANENSCH, W. 1902. Die Jurensisschichten des Elsass. *Abhandlungen zur Geologischen Specialkarte von Elsass-Lothringen; Neue Folge*, **5**, 151 pp., 12 pls.
- JAWORSKI, E. 1926. La Fauna del Lías y Dogger de la Cordillera Argentina. *Parte Meridional de la Provincia de Mendoza, III, Actas de la Academia Nacional de Ciencias*, 139–317, pls 1–4.
- KALACHEVA, E. D. and SEY, I. I. 1970. *Tugurites* – a new late Aalenian north-Pacific genus. *Doklady Akademii Nauk SSR, Seriya Geologiya*, **193**, 449–452.

- KNITTER, H. and OHMERT, W. 1983. Das Toaricum an der Schwärze bei Badenweiler (Oberrheingebiet S Freiburg). *Jahrbuch des Geologischen Landesamt Baden-Württemberg*, **25**, 233–281, pls 1–5.
- KOBAYASHI, T. 1935. Contributions to the Jurassic Torinosu Series of Japan. *Japanese Journal of Geology and Geography*, **12**, 69–91, pls 12–13.
- KOVÁCS, L. 1939. Bemerkungen zur systematischen Einteilung der jurassischen Phylloceraten. *Abhandlung Mineralogische und Geologische Institut Tisza-Universite*, **13**, 278–320.
- MATSUMOTO, T. and ONO, A. 1947. A biostratigraphic study of the Jurassic Toyora Group, with special reference to ammonites. *Scientific Reports of the Faculty of Science, Kyushu University, Geology*, **2**, 20–31, pls 1–2.
- MAUBEUGE, P. L. 1949. Revision des ammonites du genre *Phlyseogrammoceras*. *Bulletin de l'Institut Royal des Sciences Naturelles de Belgique*, **25**, 1–17, pls 1–2.
- 1950. Nouvelles recherches stratigraphiques et paléontologiques sur l'Aalénien Luxembourgeois. *Archives Institut Grand-Ducal de Luxembourg, Section des Sciences naturelles, physiques et mathématiques, nouvelle série*, **19**, 365–397, 16 pls.
- MOERICKE, W. 1894. Versteinerungen des Lias und Unteroolith von Chile. *Neues Jahrbuch für Mineralogie, Geologie und Palaeontologie, Beilage*, **9**, 1–100, pls 1–6.
- MONESTIER, J. 1931. Ammonites rares ou peu connues et ammonites nouvelles du Toarcien Moyen de la région sud-est de l'Aveyron. *Mémoires de la Société Géologique de France (Nouvelle Série)*, **15**, 5–79, pls 1–9.
- NEUMAYR, M. 1871. Jurastudien. 3. Die Phylloceraten des Dogger und Malm. *Jahrbuch von der kaiserlichen und königlichen geologischen Reichsanstalt*, **21**, 297–354, pls 12–17.
- 1875. Die Ammoniten der Kreide und die Systematik der Ammonitiden. *Zeitschrift der Deutschen geologischen Gesellschaft*, **27**, 854–942.
- O'BRIEN, J. A. 1987. Jurassic biostratigraphy and evolution of the Methow Trough, southwestern British Columbia. Unpublished M.Sc. thesis, University of Arizona, 149 pp.
- ORBIGNY, A. d'. 1840–1842. *Paléontologie française. Terrains crétacés, I: Céphalopodes*. Paris, 662 pp., 148 pls.
- 1842–1851. *Paléontologie française. Terrains jurassiques, I: Céphalopodes*. Paris, 642 pp., 234 pls.
- PARISCH, C. and VIALE, C. 1906. Contribuzione alla studio della ammoniti del Lias superiore. *Rivista Italiana di Paleontologia e Stratigrafia*, **12**, 141–168, pls 7–11.
- POULTON, T. P. and TIPPER, H. W. 1991. Aalenian ammonites and strata of western Canada. *Bulletin of the Geological Survey of Canada*, **411**, 71 pp., 7 pls.
- PRINZ, G. 1904. Die Fauna der älteren Jurabildungen im nordöstlichen Bakony. *Mitteilungen aus dem Jahrbuch der Königlichen Ungarischen Geologischen Anstalt*, **15**, pls 1–40.
- QUENSTEDT, F. A. 1883–1888. *Die Ammoniten des schwäbischen Jura*. Schweizerbart, Stuttgart, 3 vols, 1140 pp., 126 pls.
- REINECKE, I. C. M. 1818. *Nautilus et Argonautas vulgo Cornua Ammonis in agro Coburgico et vicino reperiundos*. Coburg, 90 pp., 13 pls.
- ROMAN, F. 1938. *Les Ammonites jurassiques et créteécées. Essai de genera*. Masson, Paris, 554 pp., 53 pls.
- SCHINDEWOLF, O. H. 1964. Studien zur Stammesgeschichte Ammoniten. *Akademie der Wissenschaften und der Literatur, Mainz*, **1963**(6), 287–432.
- SCHLEGELMILCH, R. S. 1976. *Die ammoniten des süddeutschen Lias*. Gustav, Fischer, Verlag, Stuttgart, 212 pp., 48 pls.
- 1985. *Die Ammoniten des süddeutschen Doggers*. Gustav, Fischer, Verlag, Stuttgart, 284 pp., 59 pls.
- SETTI, F. L. 1968. Ammoniti del genere *Dumortieria* nella serie Toarciana del Passo del Furlo (Appennino Centrale). *Atti della Società Italiana di Scienze Naturali e del Museo Civico di Storia Naturale di Milano*, **107**, 317–347, pls 30–32.
- SIMPSON, M. 1843. *A monograph of the ammonites of the Yorkshire Lias*. London, 60 pp.
- 1855. *The fossils of the Yorkshire Lias; described from nature*. London and Whitby, 149 pp.
- SMITH, P. L. 1980. Correlation of the members of the Jurassic Snowshoe Formation in the Izee basin of east-central Oregon. *Canadian Journal of Earth Sciences*, **17**, 1603–1608.
- 1986. The implications of data base management systems to paleontology: a discussion of Jurassic ammonoid data. *Journal of Paleontology*, **60**, 327–340.
- and TIPPER, H. W. 1986. Plate tectonics and paleobiogeography: Early Jurassic (Pliensbachian) endemism and diversity. *Palaios*, **1**, 399–412, pl. 1.
- SOUTHER, J. G. 1971. Geology and mineral deposits of Tulsequah map-area, British Columbia. *Memoir of the Geological Survey of Canada*, **362**, 84 pp.
- SPATH, L. F. 1927. Revision of the Jurassic cephalopod faunas of Kachh (Cutch). *Palaeontographica Indica, new series*, **9**(2), parts 1–6, 945 pp., 130 pls.

- SUTHERLAND, BROWN, A. 1968. Geology of the Queen Charlotte Islands, British Columbia. *Bulletin of the British Columbia Department of Mines and Petroleum Resources*, **54**, 226 pp.
- TAYLOR, D. G., CALLOMON, J. H., HALL, R., SMITH, P. L., TIPPER, H. W. and WESTERMANN, G. E. G. 1984. Jurassic ammonite biogeography of western North America: the tectonic implications. 121–141. In WESTERMANN, G. E. G. (ed.). Jurassic-Cretaceous biochronology and biogeography of North America. *Geological Society of Canada Special Paper*, **27**, 315 pp.
- THÉOBALD, N. 1950. Paléogéographie du jurassique supérieur dans le sud-ouest de l'Allemagne et la nord-ouest de la France. *Naturforschende Gesellschaft zu Freiburg im Breisgau, Berichte*, **40**, 123–146.
- and MOINE, H. 1959. Les Ammonites du Toarcian Supérieur et de l'Aalenien du Sentier de l'Ehn près d'Obernai (Bas-Rhin). *Bulletin du Service de la Carte Géologique d'Alsace et de Lorraine*, **12**, 3–36, pls 1–4.
- THOMSON, R. C., SMITH, P. L. and TIPPER, H. W. 1986. Lower to Middle Jurassic (Pliensbachian to Bajocian) stratigraphy of the northern Spatsizi area, north-central British Columbia. *Canadian Journal of Earth Sciences*, **23**, 1963–1973.
- TIPPER, H. W. and RICHARDS, T. A. 1976. Jurassic stratigraphy and history of north-central British Columbia. *Bulletin of the Geological Survey of Canada*, **270**, 73 pp.
- SMITH, P. L., CAMERON, B. E. B., CARTER, E. S., JAKOBS, G. K. and JOHNS, M. J. 1991. Biostratigraphy of the Lower Jurassic formations of the Queen Charlotte Islands, British Columbia. 203–235. In WOODSWORTH, G. J. (ed.). Evolution and hydrocarbon potential of the Queen Charlotte Basin, British Columbia, *Geological Survey of Canada Paper*, **90-10**, 569 pp.
- UMHOFER, P. J. 1989. Stratigraphy and tectonic setting of the Upper Cadwallader Terrane and overlying Relay Mountain Group, and the Cretaceous to Eocene Structural Evolution of the Eastern Tyaughton Basin, British Columbia. Unpublished Ph.D. thesis, University of Washington, 186 pp.
- VILLALTA, J. F. de and ROSELL, S. J. 1966. Estudio de un afloramiento Liasico en el prepirineo de la Provincia de Barcelona. *Notas y Comunicaciones, Instituto Geologico y Minero España*, **88**, 7–14, 1 pl.
- WHITE, C. A. 1889. Mesozoic molluscs from the southern coast of the Alaskan Peninsula. *Bulletin of the United States Geological Survey*, **51**, 64–70, pls 12–14.
- WHITEAVES, J. F. 1889. Fossils from the Rocky Mountains three miles north of the east end of Devil's Lake. *Geological Survey of Canada, Contributions to Canadian Palaeontology*, **1**(2), 163–172.
- WUNSTORF, W. 1907. Die Fauna der Schichten mit *Harpoceras dispansum* Lycett vom Gallberg bei Salzgitter. *Jahrbuch der Königlichen Preussischen Geologischen Landesanstalt zu Berlin für das Jahr*, **25**(3), 488–525, pls 17–20.
- YOUNG, G. M. and BIRD, J. 1828. *A geological survey of the Yorkshire Coast: describing the strata and fossils occurring between the Humber and the Tees, from the German ocean to the plain of York*, 2nd edition. Whitby, and London, 368 pp., 17 pls.
- ZIETEN, C. H. 1830–1833. *Die Versteinerungen Württembergs*. Stuttgart, 102 pp., 72 pls.
- ZITTEL, K. A. von 1884. *Handbuch der Palaeontologie*, **1**(2). Munich and Leipzig, 893 pp.

GISELLE K. JAKOBS

4714 Halliwell Avenue Terrace
British Columbia, V6G 2J3
Canada

PAUL L. SMITH

Department of Geological Sciences
University of British Columbia
6339 Stores Road, Vancouver
British Columbia, V6T 1Z4
Canada

Typescript received 29 November 1993

Revised typescript received 31 March 1995